



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

ASIAN JOURNAL OF
SCIENCE AND TECHNOLOGY

Asian Journal of Science and Technology
Vol.07, Issue, 04, pp.2753-2755, April, 2016

RESEARCH ARTICLE

SEASONAL METABOLIC VARIATION OF LIPID CONTENT IN FRESHWATER BIVALVE *LAMELLIDENS MARGINALIS* (LAMARCK, 1819)

*Padewar, S.K.

Department of Zoology, Yeshwant Mahavidyalaya, Nanded

ARTICLE INFO

Article History:

Received 26th January, 2015
Received in revised form
18th February, 2016
Accepted 07th March, 2016
Published online 27th April, 2016

Key words:

Lipid, *Lamellidens marginalis*,
Seasonal.

ABSTRACT

Lipid is a water-insoluble biomolecule. Lipid has a high solubility in nonpolar organic solvents such as chloroform. The simplest lipids are the fats. The term fats is also used as a general synonym for lipids, so the more precise terms triacylglycerols or triglycerides are preferable for the simplest lipids. Triacylglycerols are used primarily for energy storage in animals. More complex lipids are phospholipids, glycolipids, and cholesterol, are the major constituents of biological cell membranes. All different body tissues shows increasing order of Lipid contents in rainy season, minimum amount of lipid values in winter season and more amount of lipid in the summer season.

Copyright © 2016, Padewar. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

A lipid is defined as a water-insoluble biomolecule which has a high solubility in nonpolar organic solvents such as chloroform. The simplest lipids are the fats, which are triesters made up of one glycerol and three fatty acids. The term fats is also used as a general synonym for lipids, so the more precise terms triacylglycerols or triglycerides are preferable for the simplest lipids. Triacylglycerols are used primarily for energy storage in animals. More complex lipids, the phospholipids, glycolipids, and cholesterol, are the major constituents of biological cell membranes. Seasonal changes in biochemical composition have been reported by many workers. Ansell et al. (1964), De Zwann and Zandee (1972), Gabbott and Bayne (1973) determined seasonal changes in biochemical composition of adductor muscle, mantle, siphon and foot in *Mercenaria mercenaria* and *Mytilus edulis*. From India, relatively very few investigators such as Bidarkar (1975) on *Crassostrea cucullata*, Dhamane (1975) on *Paphia laterisulca*, Nagabhushanam and Mane (1975, 1978) on *Mytilus viridis*, have reported changes in the biochemical composition correlating with annual reproductive cycle of bivalves. Lipids are responsible for variety of functions in molluscs' Lipid composition in different tissues has been reported by Trumen and Pekkarinen (1990) in *Macoma balthica*.

Vedpathak (1989) observed fortnightly and monthly changes in biochemical composition in freshwater molluscs. The review of above literature shows that there is no adequate information about freshwater mollusc, *Lamellidens corrianus* from different rivers of Maharashtra. Hence, the present study was carried out to understand the changes in biochemical composition through collection of animals from Godavari river near Nanded.

MATERIALS AND METHODS

Bivalve *Lamellidens corrianus* samples (75-80 mm in shell length) were obtained from fishermen's catch. In the present study they were collected from Godavari river near district Nanded of Maharashtra in India. Immediately after bringing to laboratory, hard shells of these freshwater bivalves were brushed and washed with fresh and clean water to remove algal biomass, mud and other waste material. The cleaned animals were then kept for depuration for 14hrs in laboratory conditions under constant aeration. For Biochemical analysis, animals were dissected and soft body tissues like mantle, hepatopancreas, gonad, Adductor muscle and gill were removed. 100 mg of each wet tissues were taken for biochemical analysis. Lipids was estimated by using Methanol-Chloroform Method (Raymont et.al.,1964). The values of lipids were expressed in terms of Lipids in mg/g dry weight tissue.

*Corresponding author: Padewar, S.K.

Department of Zoology, Yeshwant Mahavidyalaya, Nanded

RESULTS AND DISCUSSION

Lipid content in *Lamellidens marginalis*

1) Mantle :- The seasonal change in the lipid content in mantle of *L.marginalis* are shown in (Table and Figure). The percentage of lipid was found to be maximum in summer and varies from 11.0325 ± 0.5236 to 12.1342 ± 0.4213 , increasing in rainy season and varies from 5.3810 ± 0.3203 to 10.1322 ± 0.6330 , whereas it is minimum in winter season and varies from 4.0432 ± 0.3512 to 6.2104 ± 0.5408 .

2) Gonad :- The seasonal change in the lipid content in gonad of *L.marginalis* are shown in (Table and Figure). The percentage of lipid was found to be maximum in summer and varies from 10.1221 ± 0.7200 to 13.1332 ± 0.3212 , increasing in rainy season and varies from 5.4230 ± 0.7123 to 9.2325 ± 0.4821 , whereas it is minimum in winter season and varies from 5.1417 ± 0.3109 to 7.4205 ± 0.1003 .

3) Hepatopancreas :- The seasonal change in the lipid content in hepatopancreas of *L.marginalis* are shown in (Table and Figure). The percentage of lipid was found to be maximum in summer and varies from 8.3310 ± 0.3210 to 11.1000 ± 0.3243 , increasing in rainy season and varies from 5.4310 ± 0.4232 to 10.2301 ± 0.4203 , whereas it is minimum in winter season and varies from 5.0201 ± 0.4101 to 6.3510 ± 0.3241 .

4) Add.Muscle:- The seasonal change in the lipid content in Add.muscle of *L.marginalis* are shown in (Table and Figure). The percentage of lipid was found to be maximum in summer and varies from 9.4023 ± 0.3220 to 11.5201 ± 0.4152 , increasing in rainy season and varies from 8.2100 ± 0.2021 to 10.1302 ± 0.3022 , whereas it is minimum in winter season and varies from 3.3401 ± 0.3000 to 5.1201 ± 0.4320 .

5) Gill :- The seasonal change in the lipid content in gill of *L.marginalis* are shown in (Table and Figure). The percentage of lipid was found to be maximum in summer and varies from 7.1010 ± 0.5210 to 10.4010 ± 0.1300 , increasing in rainy season and varies from 8.4102 ± 0.2001 to 10.1203 ± 0.5100 , whereas it is minimum in winter season and varies from 5.0200 ± 0.3231 to 6.3210 ± 0.4046 .

DISCUSSION

The investigation shows that, there is significant variation in the biochemical composition in different body tissues according to seasonal changes. Organic constituents like protein, glycogen and lipids act as key substances for different metabolic activities. Seasonal changes in lipid content is most important in relation to energy metabolism necessary for growth and reproduction (Jayabal and Kalyani, 1986; Navarro et al., 1989; Lodeiros et al., 2001). All different body tissues show increasing order of Lipid contents in rainy season, which is correlated with highest body activities of animal during this season. And due to increase inflow and turbidity of water and to cope up with the new environmental changes. It might be due to favourable environmental conditions, the period of growth with the gonadal development and lots of food availability. Similar conclusions were reported in *M. edulis*, in Britishwater by Williams, 1969 and Mane and Nagabhusanam

and Mane 1978. Lipid variation is related to gamete development (Martinez, 1991) with the highest huge levels of lipids during the period when gonads are ripe. All the body organs show minimum amount of lipid values in winter season, which may be due to sedentary life without much body activities. Lipid is an important dietary constituent, serve as reserve energy when food supply is scanty. In stressful environmental conditions, after glycogen lipid is use as energy source (Shigmatas and Takeshita, 1959; Chourpagar and Kulkarni, 2011). It is observed that lipid contents are significantly accumulated is found to be more during the summer season in all body organs due to exposure to high temperature in summer season. Similar results are observed by Pandit (2005) by *Lamellidens marginalis* of Godavari river at Kaigaon due to exposure of mantle and foot to high temperature. The study revealed that in term of energy conservation. The organic would be exported to make compensatory adjustments to both the components of energy gain and energy loss fate of changes in the environmental conditions (Vedpathak, 1989). The relationship of the energy transfer between different tissues, their capacity of reserve amounts under food availability, and their positive relationship with the high temperature and gonadal maturation have been shown in different species of bivalve molluscs such as scallops (Robinson et al., 1981; Sundet and Vahl, 1981; MacDonald and Thompson, 1986; Villalaz, 1994), mussels (Zandee et al., 1980) and clams (Robert et al., 1993; Urrutia et al., 2001). Thus, in the present study organic constituents present in different body tissues shows seasonal changes and are correlated with the change in environmental conditions along with development.




Fig: Monthly variation in lipid content of *Lamellidens marginalis*

REFERENCES

- Bayne, B. L. and Thompson, R. T. 1970. Some physiological consequences of keeping *Mytilus edulis* in the Laboratory. *Helgolander Wissenschaftliche Meeresunter Suchungen*. 20: 528-552.
- Chourpagar, A. R. and Kulkarni, G. K. 2011. Effect of heavy metal pesticides on biochemical constituents in freshwater crab *Barytelphusa cunicularis* (westwood). *Proc. Zool. Soc. India*. 10(2): 21-30
- MacDonald, B. A. and Thompson, R. J. 1986. Influence of temperature and food availability on the ecological energetics of the giant scallop *Placopecten magellanicus*. *Mar. Biol.* 93: 37-48.

- Martinez, G. 1991. Seasonal variations in biochemical composition of three size classes of the Chilean scallop, *Argopecten purpuratus* Lamark. 17: 113-116
- Nagabhushanam, R. and Mane, U. H. 1978. Seasonal variations in the biochemical composition of *Mytilus viridis* at Ratnagiri on the West Coast of India. *Hydrobiol.* 57: 69 – 72.
- Pandit, SV, 2005. Seasonal variation in the biochemical of the freshwater bivalve mollusks, *Lamellidens marginalis*(L) from Godavari river at Kaigaon, near Aurangabad. *Ph.D Thesis Dr. Babasahb Ambedkar Marathwada University, Aurangabad, India* pp. 1-213.
- Robert, R., Trut, G. and Laborde, J. L. 1993. Growth, reproduction and gross biochemical composition of the Manila clam *Ruditapes philippinarum* in the Bay of Arcachon, France. *Mar. Biol.* 116: 291- 299.
- Robinson, W. E., Wehling, W. E., Morse, M. P. and McLeod, G. C. 1981. Seasonal changes in soft-body component indices and energy reserves in the atlantic deep-sea scallop.
- Shigmates, H. and Takeshita, H. 1959. On the changes in the weight of fat body and its chief constituents in the silkworm, *Bombyx mori* L. during metamorphosis. *Appl. Entm. Zool. Japan.* 3: 123-126.
- Sundet, J. H. and Vahl, O. 1981. Seasonal changes in dry weight and biochemical composition of the tissues of sexually mature and immature Iceland scallops, *Chlamys islandica*. *J. Mar. Biol. Assoc. U.K.* 61: 1001-1010
- Urrutia, G. X., Navarro, J. M., Closing, E. and Stead R. A. 2001. The effects of environmental factors on the biochemical composition of the bivalve *Tagelus dombeii* (Lamarck, 1818) (Tellinacea: Solccurtidae) from the intertidal flat of Coihuin, Puerto Montt, Chile. *J. Shellfish Res.* 20: 1077-1087.
- Vedpathak, A. N. 1989. Reproductive endocrinology of some *Lamellibranch* molluscs with special reference to environmental stress. Ph. D. Thesis, Marathwada University, Aurangabad (M.S.), India. pp. 1 – 280.
- Villalaz, G. J. R. 1994. Morphometric and biochemical changes in two age classes of the tropical scallop, *Argopecten ventriosus*, under laboratory conditions. *Am. Mal. Bull.* 11: 67-72.
- Zandee, D. I., Kluytmans, J. H. and Zurburg, W. 1980. Seasonal variations in biochemical composition of *Mytilus edulis* with reference to energy metabolism and gametogenesis. *Neth. J. Sea Res.* 14:1-29
