



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

ASIAN JOURNAL OF
SCIENCE AND TECHNOLOGY

Asian Journal of Science and Technology
Vol. 07, Issue, 11, pp.3785-3787, November, 2016

RESEARCH ARTICLE

ON SPECIAL D(2)-QUADRUPLE

*Gopalan, M.A., Vidhyalakshmi, S. and Shanthy, J.

Department of Mathematics, SIGC, Trichy-620002, Tamilnadu

ARTICLE INFO

Article History:

Received 24th August, 2016
Received in revised form
16th September, 2016
Accepted 17th October, 2016
Published online 30th November, 2016

ABSTRACT

This paper concerns with the study of constructing a special Dio quadruple (a,b,c,d) such that the product of any two elements of the set difference with their sum and increased by 2 is a perfect square.

Key words:

Diophantine Quadruples,
Pell equation 2010 Mathematics
Subject classification 11D99).

Copyright © 2016, Thomas M. Kollars. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

The problem of constructing the sets with property that the product of any two of its distinct elements is one less than a square has a very long history and such sets were studied by Diophantus[1]. A set of m positive integers $\{a_1, a_2, \dots, a_m\}$ is said to have the property D(n), $n \in \mathbb{Z} \setminus \{0\}$ if $a_i a_j + n$, a perfect square for all $1 \leq i < j \leq m$ and such a set is called a Diophantine m-tuples with property D(n). Many mathematicians considered the construction of different formulations of Diophantine quadruples with the property D(n) for any arbitrary integer n and also for any linear polynomials in n. In this context, one may refer [2-16] for an extensive review of various problems on Diophantine quadruples. This paper aims at constructing special dio – quadruple where the special mention is provided because it differs from the earlier one and the special Dio – quadruple is constructed where the product of any two members of the quadruple with the difference of the sum of the same members and the addition of seventeen satisfies the required property.

METHOD OF ANALYSIS

Let $a(k, n) = \frac{(k+1)}{2\sqrt{k^2-1}} g(k, n) - 1$ and $b(k, n) = \frac{(k-1)}{2\sqrt{k^2-1}} g(k, n) - 1$, where

$g(k, n) = \left[(k + \sqrt{k^2 + 1})^{n+1} - (k - \sqrt{k^2 + 1})^{n+1} \right]$ be two integers such that $a(k, n)b(k, n) - [a(k, n) + b(k, n)] + 2$ is a perfect square.

Let $c_N(k, n)$ be any non-zero integer such that

*Corresponding author: Gopalan, M.A.,
Department of Mathematics, SIGC, Trichy-620002, Tamilnadu

$$[a(k,n)+1]c_N(k,n)+a(k,n)+2 = p_N^2(k,n) \dots\dots\dots(1)$$

$$[b(k,n)+1]c_N(k,n)+b(k,n)+2 = q_N^2(k,n) \dots\dots\dots(2)$$

Eliminating $c_N(k,n)$ between (1) and (2), we obtain

$$[b(k,n)+1]p_N^2(k,n)-[a(k,n)+1]q_N^2(k,n)=b(k,n)-a(k,n) \dots\dots\dots(3)$$

Taking

$$p_N(k,n) = X_N(k,n) + [a(k,n)+1]T_N(k,n) \dots\dots\dots(4)$$

$$q_N(k,n) = X_N(k,n) + [a(k,n)+1]T_N(k,n) \dots\dots\dots(5)$$

in (3), we have

$$X_N^2(k,n) = [b(k,n)+1][a(k,n)+1]T_N^2(k,n) + 1 \dots\dots\dots(6)$$

which is satisfied by

$$X_N(k,n) = \frac{1}{2}F(N,k,n) \dots\dots\dots(7)$$

$$T_N(k,n) = \frac{1}{2\sqrt{[a(k,n)+1][b(k,n)+1]}}G(N,k,n)$$

Where

$$F(N,k,n) = \left[\frac{f(k,n)}{2} + \sqrt{[a(k,n)+1][b(k,n)+1]} \right]^{N+1} + G(N,k,n) = \left[\frac{f(k,n)}{2} + \sqrt{[a(k,n)+1][b(k,n)+1]} \right]^{N+1} - \left[\frac{f(k,n)}{2} - \sqrt{[a(k,n)+1][b(k,n)+1]} \right]^{N+1}$$

& $\left[\frac{f(k,n)}{2} - \sqrt{[a(k,n)+1][b(k,n)+1]} \right]^{N+1}$

in which $f(k,n) = \left[(k + \sqrt{k^2 + 1})^{n+1} + (k - \sqrt{k^2 + 1})^{n+1} \right]$

Substituting N=0,1,2 in (7) and using (4) & (1) in turn, we have

$$c_0(k,n) = \frac{k}{\sqrt{k^2 - 1}} g(k,n) + f(k,n) - 1$$

$$c_1(k,n) = [a(k,n)+1][f(n,k) + 2b(k,n) + 2]^2 + 2[f(k,n) + 2b(k,n) + 2] - 1$$

$$c_2(k,n) = [a(k,n)+1]M^2 + f(k,n)M + b(k,n)$$

where

$$M = 4[a(k,n)+1][b(k,n)+1] + 3 + 2f(k,n)[b(k,n)+1] \text{ and so on.}$$

It is seen that the quadruple $(a(k), b(k), c_{N-1}(k), c_N(k))$, N=1,2,3... is a special dio-quadruple with property D(2). A few illustrations are given below.

Table 1.

(k,n)	$(a(k,n), b(k,n), c_0(k,n), c_1(k))$	$(a(k), b(k), c_1(k), c_2(k))$
2,1	(12,3,29,6335)	(12,3,6335,1360802)
3,2	(140,69,406,16109079)	(140,69,16109079,6.368035987x10 ¹¹)
4,3	(2480,1487,7809,1.153229063x10 ¹¹)	(2480,1487,1.153229063x10 ¹¹ , 1.703149655x10 ¹⁸)
5,1	(50,39,197,1616239)	(50,39,1616239,1.306246479x10 ¹⁰)

Conclusion

This paper concerns with the construction of special dio - quadruples with property D(2). One may search for special dio-quadruples consisting of special numbers with suitable property.

REFERENCES

- Abu Muriefah F.S and Al-Rashed .A, some Diophantine quadruples in the ring $Z[\sqrt{2}]$ Math. Commun,2004,9,1-8,.
 Assaf.E and Gueron.S, Characterization of regular Diophantine quadruples, Elem. Math. 2001,56, 71-81,
 Bashmakova.I.G(ed.), Diophantus of Alexandria, Arithmetic's and the book of polynomial numbers, Nauka Moscow, 1974
 Dujella.A, Some polynomial formulas for Diophantine quadruples, Grazer Math. Ber,1996,328, 25-30.
 Dujella.A, Some estimates of the number of Diophantine quadruples, Publ. Math. Debrecen 1998,53,177-189.
 Filipin.A and Fujita. Y, Any polynomial D(4)-quadruple is regular, Math.Commun,2008,1(3) 45-55.
 Filipin.A, An irregular D(4)-quadruple cannot be extended to a quintuple,ActaArith,2009,136,167-176.
 Franusic Z., Diophantine quadruples in the ring $Z[\sqrt{2}]$ Math,Commun,2004,9,141-148.
 Franusic.Z, Diophantine quadruples in Quadratic Fields, Dissertation,University of Zagreb, 2005 (in Croatian)
 Gopalan.M.A, Srividhya.G, Diophantine quadruples for Fibonacci and Lucas Numbers with property D(4), diophantus J.Math,2012, 1 ,15-18.
 Gopalan M.A., Vidhyalakshmi. S, Thiruniraiselvi. N. A special Diophantine quadruple with property D(17),IJMTT, 2015,Vol.20(2),110-112
 Gopalan M.A, Vidhyalakshmi .S, Shanthi.J, On special D(17)-quadruple, *International Journal of Statistics and Applied Mathematics*, 2016, 1(2),04-05.
 Gibbs.P, Diophantine quadruples and Cayley's hyperdeterminant, XXX Mathematics Archive math. NT/0107203
 Gibbs.P, Adjugates of Diophantine Quadruples, Integers,2010, 10,201-209.
 G.Srividhya, Diophantine quadruples for Fibonacci numbers with property D(1), Indian journal of mathematics and Mathematical Sciences, 2009, 5,57-59.
 Vidhyalakshmi. S., Gopalan M.A., Kavitha. A. Special dio-quadruple with property D(5), IRJMEIT, Aug-2014, Vol.1(4),16-20.
