



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

ASIAN JOURNAL OF
SCIENCE AND TECHNOLOGY

Asian Journal of Science and Technology
Vol. 08, Issue, 09, pp.5634-5637, September, 2017

RESEARCH ARTICLE

SYNTHESIS AND CHARACTERIZATION OF SOME TRANSITION METAL COMPLEXES USING BIS-(α -BENZIL)-2-AMINOTHIOPHENOL

*Anita Kumari

Research Scholar, University Department of Chemistry, LNMU-DBG, India

ARTICLE INFO

Article History:

Received 08th June, 2017

Received in revised form

14th July, 2017

Accepted 30th August, 2017

Published online 27th September, 2017

Key words:

Binuclear complexes,
Bis-(α -benzil)-2-aminothiophenol,
Diketones, Molar conductance,
Octahedral.

ABSTRACT

Binuclear complexes of Co(II), Ni(II), Cu(II) and Zn(II) with bis-(α -benzil)-2-aminothiophenol, Schiff base ligand, have been synthesized and characterized on the basis of molar mass, elemental analysis, IR and electronic spectral studies, molar conductance and magnetic susceptibility measurement. On the basis of above physicochemical and spectrometric measurements it is proposed that the compounds act in a bi-dentate manner. Such complexes have varieties of useful pharmaceutical activities and many of them gained wide acceptance in clinical practice. The resulting complexes have been tested for their antifungal activities against various organisms. Complexes, excepting that of Zn(II), are colored. Electronic spectra and magnetic susceptibility study proposes octahedral geometry of the complexes.

Copyright©2017, Anita Kumari. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

An overview of the research works, going on in the field of coordination chemistry, presented by Cambridge Structural Data (CSD) base shows that the Transition Metal Chemistry has become the pioneer chemistry today. There are enumerable compounds formed by transition metal having diversified role in several aspects of our life. Several transition and inner-transition metal complexes having nitrogen and oxygen donor atoms play an important role in biological systems (Frausto da Silva, 1991; Kaim, 1996; Aayed and Al Shihri, 2004). Some of the metal complexes show carcinostatic and tuberculostatic activity (Ng *et al.*, 1953; Ma and Tien, 1953). They have industrial, antifungal, antibacterial, antiviral and herbicidal applications also (Singh *et al.*, 2007; Cozzi *et al.*, 2004; Canpolat *et al.*, 2004). There are several fascinating examples of transition metal complexes having catalytic, biological and pharmaceutical importance (Comprehensive Coordination Chemistry I, 1987; Raux *et al.*, 2000; Magnus Gustafsson *et al.*, 2010). There has been a recent upsurge in the research activity of zinc due to its recently discovered zinc finger protein and zinc complexes with scorpionate ligands (Comprehensive Coordination Chemistry II, 2005). Present programme deals with synthesis, characterization and study of a series of binuclear complexes

(Chuanbo Gao *et al.*, 2011; Mei Zhang *et al.*, 2012) of Co(II), Ni(II), Cu(II) and Zn(II) with bis-(α -benzil)-2-aminothiophenol ligand having nitrogen and sulphur donors. The complexes have been synthesized by refluxing the ethanolic solutions of metal chloride with that of bis-(α -benzil)-2-aminothiophenol with aqueous NaOH. The principal objective of the present work is to prepare some binuclear complexes having donor atoms and their characterization by means of analytical, spectral and magnetic studies. Magnetic moment data and electronic spectral bands indicate an octahedral environment around each metal ion.

Experimental

Chemicals namely bis-(α -benzil), 2-aminothiophenol (Fluka) and all solvents used in this study were obtained from Merck and used as supplied without further purification. Chlorides of transition metal were also used as supplied. For the estimation of metals, the complexes were first decomposed to bring them in their proper ionic solution and then they were quantitatively analyzed. Cobalt was estimated from its ionic solution by oxidizing it with few drops of conc. HNO₃ after heating and cooling. Now the oxide of cobalt was converted into nitrate. On expelling HNO₃ by heating, it was treated with excess of conc. H₂SO₄ to convert it into sulphate. Finally it was heated between 450-500^oc and weighed as CoSO₄. From the ionic solution of nickel chloride, nickel was precipitated as bis-DMG nickel(II) in ammonical medium. The precipitate was


*Corresponding author: Anita Kumari

Research Scholar, University Department of Chemistry, LNMU-DBG, India.

heated at 110-120°C and weighed as nickel compound. Copper was estimated by repeated evaporation of the weighed amount of its complex with HNO₃ and HClO₄ in a Pyrex beaker until a clear solution was obtained. The solution was further boiled with HCl and the volume was reduced to 10-15ml. Now the resulting solution was diluted with water and Cu was determined iodimetrically. On the other hand zinc was estimated by EDTA titration in an alkaline medium having pH 10. Elemental analysis of C, H, N and S was carried out micro-analytically at CDRI Lucknow. The electronic absorption spectra of the complexes were recorded with Hitachi-300 spectrophotometer at CDRI Lucknow. Infrared spectra were also recorded at CDRI Lucknow on Perkin Elmer 577 spectrophotometer using KBr disc in the range of 4000-400 cm⁻¹. Magnetic susceptibility of all the complexes were measured by Guoy's method using Hg[Co(NCS)₄] as calibrant. Electrical conductivity of the solution of complexes was measured by conductivity meter systronics model 300 at room temperature in DMF using N/10 and N/100 KCl solution. Pure DMF and conductivity water were used as solvent.

Synthesis of ligand

The ligand bis-(α -benzil)-2-aminithiophenol was synthesized by refluxing an ethanolic solution of α -benzil (2.1g, 0.01mol.) with an ethanolic solution of 2-aminothiophenol (0.02 mol.) for ~ 6 hours. The resulting solution was concentrated and the precipitate was separated by filtration, washed with ethanol and then air-dried. Following is the chemical reaction taking place during synthesis of bis-(α -benzil)-2-aminithiophenol.




Synthesis of bis-(α -benzil)-2-aminithiophenol

Synthesis of complexes

Following is the general method used to synthesize the binuclear complexes of Co, Ni, Cu and Zn with bis-(α -benzil)-2-aminithiophenol. An ethanolic chloride solutions of cobalt, nickel, copper and zinc were added separately to the ethanolic solution of bis-(α -benzil)-2-aminithiophenol in a molar ratio (2 : 2 = M : L) or (1 : 1 = M : L) followed by the addition of an aqueous solution of NaOH (0.02 mol.) to the resulting mixture. Now all the mixtures were refluxed separately for about 4 hours. The formed precipitates were collected by filtration, washed several times with ethanol and water and finally dried

in oven at 110°C. The synthetic reaction during the synthesis of metal complexes can be represented as



RESULTS AND DISCUSSION

The complexes obtained from bis-(α -benzil)-2-aminithiophenol were colored solid, stable in air at room temperature. Insolubility of these complexes in common organic solvents such as acetone, methanol and chloroform along with their higher range of decomposition temperature support the binuclear structure of these complexes. The molar conductance of the complexes in DMSO was in the range of (15-7) ohm⁻¹ cm² mol⁻¹. On the basis of above data these complexes can be considered as non-electrolyte (Jun Huang *et al.*, 2007).

IR-spectral studies

The characteristic IR spectral bands of the complexes when compared with that of the free ligand provide ample evidences for the mode of binding of the ligand to the metal ion. In the IR spectrum of bis-(α -benzil)-2-aminithiophenol ligand, a sharp band observed at 1594 cm⁻¹ was assigned to the $\nu(\text{C}=\text{N})$ mode of the azomethine group which was shifted to lower wave numbers in all the complexes, suggesting the coordination of the azomethine nitrogen to the metal ion centres (Gopalakrishna Bhat and Narayana, 2005). This is further substantiated by the presence of the new band at 370-390 cm⁻¹ assignable to $\nu(\text{M}-\text{N})$. The characteristic thiophenolic $\nu(\text{S}-\text{H})$ mode was observed at 2475 cm⁻¹. A band at 1293 cm⁻¹ due to $\nu(\text{C}-\text{S})$ thiophenolic was also observed in the spectrum of ligand (Singh *et al.*, 2006). The disappearance of thiophenolic $\nu(\text{S}-\text{H})$ bond in the complexes under study suggests the coordination by thiophenolic sulphur after deprotonation to the metal ion. This is further supported by the shifting of $\nu(\text{C}-\text{S})$ thiophenolic to lower wave numbers in all the metal complexes. The IR spectra of the complexes also show a band in the 3411-3452 cm⁻¹ region, indicating the presence of coordinated water in these complexes. The presence of coordinated water was further confirmed by the appearance of a non-ligand band in the 851-892 cm⁻¹ and 744-764 cm⁻¹ regions, assignable to the rocking and wagging modes of water, respectively (Singh *et al.*, 2010).

Electronic spectra and magnetic moment studies

The electronic spectra of the free ligand, bis-(α -benzil)-2-aminithiophenol, and its Co(II) complexes were recorded in DMSO at room temperature. Electronic absorption spectral data of the ligand shows the $\pi \rightarrow \pi^*$ transitions related to benzene ring in the range 43000-44000 cm⁻¹ and imine $n \rightarrow \pi^*$ transitions in the range 30500-31000 cm⁻¹. The [(Co)₂(L)₂(H₂O)₄] complex exhibits one band in the range 9850-10000 and another band in the range of 23800-23900 cm⁻¹. The bands are assigned to the transitions ⁴T_{1g}(F) \rightarrow ⁴T_{2g}(F)(ν_1) and ⁴T_{1g}(F) \rightarrow ⁴T_{1g}(P)(ν_3) respectively (Lever, 1968). The magnetic moment of the complex, [(Co)₂(L)₂(H₂O)₄], at room temperature was observed in the range of 4.20-4.60 B.M. per Co atom.



Table 1. Colour, yield, decomposition temperature, analytical and molar conductance data of the complexes of bis-(α -benzil)-2-aminothiophenol

Compound	Colour	Yield (Gram)	Decomp. (Degree)	Analytical Data (Cal./Found %)				Cal.(Found) %		\wedge_M ohm ⁻¹ cm ² mol ⁻¹
				C	H	N	S	Metal		
[(Co) ₂ L ₂ (H ₂ O) ₄]	Dark Brown	69	327	73.57/73.02	4.72/4.69	6.61/6.59	15.09/14.97	11.41(11.37)		7
[(Ni) ₂ L ₂ (H ₂ O) ₄]	Yellow	73	355	73.57/73.02	4.72/4.69	6.61/6.59	15.09/14.97	11.42(11.38)		9
[(Cu) ₂ L ₂ (H ₂ O) ₄]	Brown	76	320	73.57/73.02	4.72/4.69	6.61/6.59	15.09/14.97	12.19(11.98)		11
[(Zn) ₂ L ₂ (H ₂ O) ₄]	Yellow	77	340	73.57/73.02	4.72/4.69	6.61/6.59	15.09/14.97	12.50(12.47)		15


Table 2. Important IR spectral data of the complexes of bis-(α -benzil)-2-aminothiophenol (cm⁻¹)

Compound	ν (H ₂ O)	ν (C=N)	Rocking ν (H ₂ O)	Wagging ν (H ₂ O)	ν (M-N)	ν (M-S)
[(Co) ₂ L ₂ (H ₂ O) ₄]	3411	1592	892	756	379	494
[(Ni) ₂ L ₂ (H ₂ O) ₄]	3414	1593	868	753	390	487
[(Cu) ₂ L ₂ (H ₂ O) ₄]	3447	1594	875	764	374	494
[(Zn) ₂ L ₂ (H ₂ O) ₄]	3452	1584	851	744	370	490

On the basis of the position of the bands and the magnetic moment value the geometry of [(Co)₂(L)₂(H₂O)₄] complex is presumably octahedral. The Ni(II) complex, [(Ni)₂(L)₂(H₂O)₄], reported herein was found to have room temperature magnetic moment value in the range of 3.75-3.80 B.M. corresponding to two unpaired electrons. Electronic spectra of the aforesaid complex display three absorption bands in the range 11250-11570, 16200-16350, and 28115-26240 cm⁻¹. These bands can be assigned to three spin allowed transitions ³A_{2g}(F) → ³T_{2g}(F), ³A_{2g}(F) → ³T_{1g}(F) and ³A_{2g}(F) → ³T_{1g}(P) respectively. The position of these bands indicates that the complex has an octahedral geometry around Ni(II) ions. The electronic spectra of Cu(II) complex, [(Cu)₂(L)₂(H₂O)₄], shows a single broad band in the region 20000-20300 cm⁻¹ attributable to ²T_{2g} → ²E_g transition indicative of distorted octahedral geometry around Cu(II) ions. A low energy band at 10341 cm⁻¹ and the magnetic moment in the range of 1.75-1.78 B.M. per Cu(II) ion support the octahedral geometry the copper complex. Zn(II) complex, [(Zn)₂(L)₂(H₂O)₄], shows zero magnetic moment suggesting the diamagnetic nature of the complex. The electronic spectra of Zn(II) complex shows a single broad band in the region 23000-25000 cm⁻¹ attributable to charge transfer transition. The electronic spectroscopy does not permit the establishment of a clear stereochemistry for Zn(II) complex. But taking into consideration of similarity with light congeners, Zn(II) complex has been proposed to have octahedral geometry. Hence the proposed structure of transition metal complexes [(M)₂(L)₂(H₂O)₄] having octahedral geometry is shown as :

**The Proposed Structure of the complexes**

The Proposed Structure of the Complexes



[M= Co(II), Ni(II), Cu(II) or Zn(II); R = C₆H₅]

Conclusion

Present programme reveals that complexes of Co(II), Ni(II), Cu(II) and Zn(II) with bis-(α -benzil)-2-aminothiophenol assume octahedral geometry by virtue of their magnetic moment values but Zn(II) complex, diamagnetic in nature having zero magnetic moment value, is proposed to have octahedral geometry due to its similarity with light congeners.

Acknowledgment

The authors are very thankful to the Principal and P.G. Department of Chemistry, Samastipur College, Samastipur for providing all necessary facilities as well as to the Director of CDRI Lucknow for recording the infrared and electronic spectra. The micro-analysis of samples was also done by CDRI Lucknow.

REFERENCES

- Ayed, S., Al Shihri, 2004. *Spectrochimica Acta Part A*; 60 1189.
- Canpolat, E., Kaya, M. 2004. *J. Coord. Chem*; 57, 1217.
- Chuanbo Gao, Zhenda Lu and Yadong Yin, *Langmuir*, 2001. 27 :12201.
- Comprehensive Coordination Chemistry I*, Geoffrey Wilkinson and Jon A McCleverty (Editor-in-Chief) ELSEVIER (1987).
- Comprehensive Coordination Chemistry II*, Jon A McCleverty and Thomas J Meyer (Editor-in-Chief) ELSEVIER (2005).
- Cozzi, P. G. 2004. *Chem. Soc. Rev*; 33, 410.
- Frausto da silva, J. J. R. and R. J. P. Williams, 1991. "The Biological Chemistry of Elements" Clarendon Press, Oxford.
- Gopalakrishna Bhat, N. and Narayana, B. 2005. "Synthesis and Reactivity in Inorganic, Metal-Organic and Nano-Metal Chemistry," Vol. 35, no. 3, pp. 253.
- Jun Huang, Qing-Dong Huang, Ji Zhang, Li Hong Zhou, Qiang-lin Li, Kun Li, Ning Jiang, Hong-Hui Lin, Jiang Wi and Xiac-Qi Yu, 2007. *Int. J. Mol. Soc.* 8; 606.
- Kaim, B. Schwederski, 1996. *Bioinorganic Chemistry: Inorganic Elements in the Chemistry of Life*, Wiley, New York.
- Lever, A. B. P. 1968. *Inorganic Spectroscopy*, Elsevier, msterdam, the Netherlands.
- Ma, T. S. and Tien, T. M. 1953. *Antibiotics and Chemotherapy*; 3, 491.

- Magnus Gustafsson, 2010. Andreas Fischer, Andrey Ilyukhin, Mikhail Maliatrick and Per Nordblad, *Inorg. Chem*; 49 5359.
- Mei Zhang, Dong-Meixian, Na Zhang, Zhang-Lu Yu, 2012. *J. Coord. Chem*; (2012) 1837.
- Ng. Ph. Buu – Hoi, Ng. D. Xuong, Ng. H. Ham, F. Binon and Roger, 1953. *J. Chem. Soc.*, 1358.
- Raux, E., Schubert, H. L., Warren, M. J. 2000. *Cellular and Molecular Life Sciences* 57, 1880.
- Singh, K. Barwa, M. S. Tyagi, P. 2007. *Eur. J. Med. Chem*; 42 394.
- Singh, K. Kumar, Y. and Barwa, M. S. 2010. *South African Journal of Chemistry*, Vol. 63, pp. 169, 2010.
- Singh, K., D. P. Singh, M. S. Barwa, P. Tyagi and Y. Mirza, 2006. *Journal of Enzyme Inhibition and Medicinal Chemistry*, Vol. 21, no. 6, pp. 749.
