



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

**ASIAN JOURNAL OF
SCIENCE AND TECHNOLOGY**

Asian Journal of Science and Technology
Vol. 08, Issue, 11, pp.6320-6323, November, 2017

RESEARCH ARTICLE

STUDY ON SEMICONFORMAL KAEHLERIAN RECURRENT AND SYMMETRIC SPACES OF SECOND ORDER

***Rawat, K.S. and Sandeep Chauhan**

Department of Mathematics, H.N.B. Garhwal University Campus, BadshahiThaul, Tehri, Garhwal -249199, (Uttarakhand)

ARTICLE INFO ABSTRACT

Article History:

Received 22nd August, 2017
 Received in revised form 09th September, 2017
 Accepted 06th October, 2017
 Published online 10th November, 2017

Key words:

Kaehlerian Recurrent Space,
 Kaehlerian Symmetric Spaces,
 Semiconformal Curvature Tensor.

Walker (1950) studied on Ruse’s spaces of recurrent Curvature. Singh (1971) studied on Kaehlerian spaces with recurrent Bochner Curvature tensor. Ishii [8] introduced the notion of Conharmonic transformation under which a harmonic function transforms into a harmonic function. Negi and Rawat (1994) studied some bi-recurrent and bi-symmetric properties in a Kaehlerian space. Further, Rawat and Kumar [13] studied Weyl-Sasakian Projective and Weyl-Sasakian Conformal bi-recurrent and bi-symmetric spaces. In the present paper, we have studied and defined Semiconformal Kaehlerian recurrent and symmetric spaces of second order. Several Theorems also have been established and proved therein.

Copyright©2017, Rawat and Sandeep Chauhan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

An n-dimensional Kaehlerian space is a Riemannian space which admits a tensor field satisfying

$$F_i^h F_j^i = -\delta_j^h, \dots(1.1)$$

$$F_{ij} = -F_{ji}, (F_{ij} = F_i^\alpha g_{\alpha j}) \dots(1.2)$$

and

$$F_{i,j}^h = 0 \dots(1.3)$$

Where the (,) followed by an index denotes the operation of covariant differentiation with respect to the metric tensor of the Riemannian space.

The Riemannian Curvature tensor is given by

$$R_{ijk}^h = \partial_i \left\{ \begin{matrix} h \\ j \ k \end{matrix} \right\} - \partial_j \left\{ \begin{matrix} h \\ i \ k \end{matrix} \right\} + \left\{ \begin{matrix} h \\ i \ \alpha \end{matrix} \right\} \left\{ \begin{matrix} \alpha \\ j \ k \end{matrix} \right\} - \left\{ \begin{matrix} h \\ j \ \alpha \end{matrix} \right\} \left\{ \begin{matrix} \alpha \\ i \ k \end{matrix} \right\}, \dots(1.4)$$

Where g_{ij} and F_{ij} denotes real local coordinates. The Ricci – tensor and scalar curvature are respectively given by

***Corresponding author: Rawat, K.S.**

Department of Mathematics, H.N.B. Garhwal University Campus, Badshahi Thaul, Tehri, Garhwal -249199, (Uttarakhand).

$$R_{ij} = R_{aij}^a \text{ and } R = g^{ij}R_{ij} \dots\dots\dots(1.5)$$

If we define a tensor

$$S_{ij} = F_i^a R_{aj}, \dots\dots\dots 1.6$$

Then, we have

$$S_{ij} = S_{ji}, \dots\dots\dots(1.7)$$

$$F_i^a = -S_{ia} F_j^a, \dots\dots\dots(1.8)$$

And

$$F_i^a S_{jk,a} = R_{ji,k} - R_{ki,j} \dots\dots\dots(1.9)$$

It has been verified by (Yano[4]), that the metric tensor and the Ricci tensor denoted by and hybrid in i and j , Therefore, we get

$$g_{ij} = g_{sr} F_i^s F_j^r, \dots\dots\dots(1.10)$$

$$\text{and } R_{ij} = R_{sr} F_i^s F_j^r, \dots\dots\dots(1.11)$$

The Conharmonic curvature tensor, is given by

$$H_{ijk}^h = R_{ijk}^h - \frac{1}{(n-2)} (g_{ij} R_k^h - \delta_j^h R_{ik} + \delta_k^h R_{ij} - g_{ik} R_j^h) \dots(1.12)$$

and, the Semiconformal curvature tensor , is given by

$$P_{ijk}^h = - (n - 2) b C_{ijk}^h + [a + (n - 2) b] H_{ijk}^h \dots(1.13)$$

where a, b are constants not simultaneously zero and is Conformal curvature tensor. The Conformal curvature tensor , is given by.

$$C_{ijk}^h = R_{ijk}^h - \frac{1}{(n-2)} (g_{ij} R_k^h - \delta_j^h R_{ik} + \delta_k^h R_{ij} - g_{ik} R_j^h) + \frac{R}{(n-1)(n-2)} (\delta_k^h g_{ij} - \delta_j^h g_{ik}) \dots(1.14)$$

In Particular, if a = 1 and b = - , then the Semiconformal curvature tensor reduces to Conformal curvature tensor whereas for a = 1 and b = 0, such a curvature tensor reduces into Conharmonic curvature tensor.

In view of equations (1.12) and (1.14), equation (1.13) reduces to

$$P_{ijk}^h = a [R_{ijk}^h - \frac{1}{(n-2)} (g_{ij} R_k^h - \delta_j^h R_{ik} + \delta_k^h R_{ij} - g_{ik} R_j^h)] + b \frac{R}{(n-1)} (\delta_k^h g_{ij} - \delta_j^h R_{ik}) \dots (1.15)$$

Properties of Semiconformal Kaehlerian Recurrent space of second order

Definition (2.1) :A Kaehlerian space is said to Kaehlerian recurrent space of second order, if it satisfies.

$$R_{ijk,ab}^h - \lambda_{ab} R_{ijk}^h = 0, \dots(2.1)$$

For some non - zero tensor, and is called Kaehlerian Ricci-recurrent space of second order, if it satisfies the condition

$$R_{ij,ab} - \lambda_{ab} R_{ij} = 0, \dots(2.2)$$

Multiplying the above equation by, we have

$$R_{ab} - \lambda_{ab} R = 0, \dots(2.3)$$

Remark (2.1): From (2.1) and (2.2), it follows that every Kaehlerian recurrent space of second order is Ricci – recurrent of second order, but the Converse is not necessarily true.

Definition (2.2): A Kaehlerian space satisfying the condition

$$H_{ijk,ab}^h - \lambda_{ab} H_{ijk}^h = 0, \dots(2.4)$$

For some non - zero tensor, will be called Kaehlerian recurrent space with Conharmonic curvature tensor of second order.

Definition (2.3): A Kaehlerian space satisfying the condition

$$P_{ijk,ab}^h - \lambda_{ab} P_{ijk}^h = 0, \dots(2.5)$$

For some non - zero tensor, is said to be Semiconformal Kaehlerian Recurrent space of second order.

Definition (2.4) : A Kaehlerian space satisfying the Condition

$$C_{ijk,ab}^h - \lambda_{ab} C_{ijk}^h = 0, \dots(2.6)$$

For some non - zero tensor, is said to be Kaehlerian recurrent space with Conformal curvature tensor of second order.

Now, we have the following theorems:

Theorem (2.1): If a Kaehlerian space satisfies any two of the following properties

- The space is Kaehlerian recurrent space of second order,
- The space is Kaehlerian Ricci-recurrent space of second order,
- The space is Kaehlerian recurrent space with Semiconformal curvature tensor of second order,then it must also satisfy the third.

Proof : Differentiating equation (1.15) covariantly w.r. to, again differentiate the result thus obtained covariantly w.r. to, We have

$$P_{ijk,ab}^h = a [R_{ijk,ab}^h - \frac{1}{(n-2)} (g_{ij} R_{k,ab}^h - \delta_j^h R_{ik,ab} + \delta_k^h R_{ij,ab} - g_{ik} R_{j,ab}^h)] + b \frac{R_{,ab}}{(n-1)} (\delta_k^h g_{ij} - \delta_j^h g_{ik}) \dots (2.7)$$

Multiplying (1.15) by, and subtracting the result thus obtained from (2.7), we have

$$P_{ijk,ab}^h - \lambda_{ab} P_{ijk}^h = a [(R_{ijk,ab}^h - \lambda_{ab} R_{ijk}^h) - \frac{1}{(n-2)} ((R_{k,ab}^h - \lambda_{ab} R_k^h) g_{ij} - (R_{ik,ab} - \lambda_{ab} R_{ik}) \delta_j^h + (R_{ij,ab} - \lambda_{ab} R_{ij}) \delta_k^h - (R_{j,ab}^h - \lambda_{ab} R_j^h) g_{ik})] + b \frac{(R_{,ab} - \lambda_{ab} R)}{(n-1)} (\delta_k^h g_{ij} - \delta_j^h g_{ik}) \dots (2.8)$$

The statement of the above theorem follows in view of equations (2.1), (2.2), (2.3),(2.5) and (2.8).

Theorem (2.2) :If a Kaehlerian space satisfies any two of the following properties

- The space is Kaehlerian recurrent space with Conharmonic curvature tensor of second order,
- The space is Kaehlerian recurrent space with Conformal curvature tensor of second order,
- The space is Kaehlerian recurrent space with Semiconformal curvature tensor of second order, then it must also satisfy the third.

Proof: Differentiating (1.13) covariantly w.r. to, again differentiate the result thus obtained covariantly w.r. to, we have

$$P_{ijk,ab}^h = - (n - 2) b C_{ijk,ab}^h + [a + (n - 2) b] H_{ijk,ab}^h \dots (2.9)$$

Multiplying (1.13) by , and subtracting from (2.9), we have

$$P_{ijk,ab}^h - \lambda_{ab} P_{ijk}^h = - (n - 2) b (C_{ijk,ab}^h - \lambda_{ab} C_{ijk}^h) + [a + (n - 2) b] (H_{ijk,ab}^h - \lambda_{ab} H_{ijk}^h) \dots (2.10)$$

The statement of the above theorem follows in view of equations (2.4), (2.6) and (2.10).

Theorems (2.3): The necessary and sufficient condition for a Kaehlerian recurrent space with semiconformal curvature tensor of second order to be Kaehlerian recurrent space of

second order is that the space be Ricci-recurrent space of second order.

Proof : Let the Kaehlerian recurrent space with Semiconformal curvature tensor of second order be Kaehlerian recurrent space of second order, so that equations (2.1) and (2.5) are satisfied and equation (2.8), in view of equation (2.1) and (2.5) reduces to

$$-\frac{a}{n-2}[(R_{k,ab}^h - \lambda_{ab}R_k^h)g_{ij} - (R_{ik,ab} - \lambda_{ab}R_{ik})\delta_j^h + (R_{ij,ab} - \lambda_{ab}R_{ij})\delta_k^h - (R_{j,ab}^h - \lambda_{ab}R_j^h)g_{ik}] + b \frac{(R_{,ab} - \lambda_{ab}R)}{(n-1)}(\delta_k^h g_{ij} - \delta_j^h g_{ik}) = 0,$$

$$\text{Or, } a(n-1)[(R_{k,ab}^h - \lambda_{ab}R_k^h)g_{ij} - (R_{ik,ab} - \lambda_{ab}R_{ik})\delta_j^h + (R_{ij,ab} - \lambda_{ab}R_{ij})\delta_k^h - (R_{j,ab}^h - \lambda_{ab}R_j^h)g_{ik}] - (n-2)b \cdot (R_{,ab} - \lambda_{ab}R)(\delta_k^h g_{ij} - \delta_j^h g_{ik}) = 0,$$

which after further calculation and simplification shows that the space is Ricci-recurrent space of second order. Conversely, let Kaehlerian recurrent space with semiconformal curvature tensor of second order be Ricci-recurrent space of second order, so that equation (2.2) and (2.3) are satisfied and equation (2.8), in view of equations (2.2), (2.3) and (2.5), reduces to

$$R_{ijk,ab}^h - \lambda_{ab}R_{ijk}^h = 0,$$

which shows that the space is Kaehlerian recurrent space of second order.

This Completes the proof of the theorem.

Properties of Semiconformal Kaehlerian Symmetric space of second order

Definition (3.1) : A Kaehlerian space is said to be Kaehlerian symmetric space of second order, if it satisfies.

$$R_{ijk,ab}^h = 0, \text{ or, equivalently } R_{ijkl,ab} = 0, \quad \dots(3.1)$$

and will be called Kaehlerian Ricci-symmetric space of second order, if it satisfies.

$$R_{ij,ab} = 0, \quad \dots(3.2)$$

Multiplying equation (3.2) by , we have

$$R_{,ab} = 0, \quad \dots(3.3)$$

Remark (3.1): From (3.1) and (3.2), it follows that every Kaehlerian symmetric space of second order is Ricci-symmetric space of second order, but the Converse is not necessarily true.

Definition (3.2): A Kaehlerian space satisfying the condition

$$H_{ijk,ab}^h = 0, \text{ or, equivalently } H_{ijkl,ab} = 0, \quad \dots(3.4)$$

will be called a Kaehlerian symmetric space with Conharmonic curvature tensor of second order.

Definition (3.3) : A Kaehlerian space satisfying the condition

$$P_{ijk,ab}^h = 0, \text{ or, equivalently } P_{ijkl,ab} = 0, \quad \dots(3.5)$$

will be called a Semiconformal Kaehlerian symmetric space of second order.

Definition (3.4) : A Kaehlerian space satisfying the condition

$$C_{ijk,ab}^h = 0, \text{ or, equivalently } C_{ijkl,ab} = 0, \quad \dots(3.6)$$

will be called a Kaehlerian symmetric space with Conformal curvature tensor of second order.

Now, we have the following theorems :

Theorem (3.1) : If a Kaehlerian space satisfies any two of the following properties

- The space is Kaehlerian symmetric space of second order,
- The space is Kaehlerian Ricci-symmetric space of second order,
- The space is Kaehlerian symmetric space with Semiconformal curvature tensor of second order, then it must also satisfy the third.

Proof : A Kaehlerian symmetric space of second order satisfied the relation (3.1) and Kaehlerian Ricci-symmetric space of second order and Kaehlerian symmetric space with Semiconformal curvature tensor of second order characterized by (3.2) and (3.5) respectively.

Therefore, the statement of the above theorem follows in view of equations (3.1), (3.2), (3.5) and (2.7).

Theorem (3.2): If a Kaehlerian space satisfies any two of the following properties

- The space is Kaehlerian symmetric space with Conharmonic curvature tensor of second order,
- The space is Kaehlerians symmetric space with Conformal curvature tensor of second order,
- The space is Kaehlerian symmetric space with Semiconformal curvature tensor of second order, then it must also satisfy the third.

Proof : Kaehlerian symmetric space with Conharmonic curvature tensor of second order, Kaehlerian symmetric space with Semiconformal curvature tensor of second order and Kaehlerian symmetric space with Conformal curvature tensor of second order are characterized by (3.4), (3.5) and (3.6) respectively.

Therefore, the statement of the above theorem follows in view of equations (3.4), (3.5), (3.6) and (2.9). **Theorem (3.3) :** The necessary and sufficient condition for a Kaehlerian symmetric space with Semiconformal curvature tensor of second order to be Kaehlerian symmetric space of second order is that the space be Ricci symmetric space of second order. **Proof :** Kaehlerian symmetric space of second order and Kaehlerian symmetric space with Semi-conformal curvature tensor of

second order characterized by the equation (3.1) and (3.5) respectively.

The statement of the above theorem follows in view of equations (3.1), (3.2), (3.3), (3.5) and (2.7).

Conversely, If Ricci symmetric space of second order and Kaehlerian symmetric space with Semi-conformal curvature tensor of second order given by the equations (3.2) and (3.5) respectively. Therefore, by using equations (3.2), (3.3) and (3.5) in the equation (2.7), we have

$$R_{ijk,ab}^h = 0,$$

which shows that the space is Kaehlerian symmetric space of second order.

REFERENCES

- Ishii, Y. 1957. On Conharmonic transformations, *Tensor (N.S)* 7, 73-80.
- Negi, D. S. and Rawat, K. S. 1994. Some bi-recurrent and bi-symmetric properties in a Kaehlerian space, *Acta Ciencia Indica*, Vol. XXM, No.1, 95-100.
- Okumura, M. 1962. Some Remarks on space with a certain contact structures, *Tohoku Math. Jour.*, Vol.14, PP 135-145.
- Otsuki, T. and Tashiro, Y. 1954. On Curves in Kaehlerian spaces. *Math. J. Okayama University*, Vol. 4, PP 57-78.
- Rawat, K. S. and Kunwar Singh 2012. Some investigations in Kaehlerian Tachibana H-Concircular recurrent and symmetric spaces of second order, *Acta Ciencia Indica*, Vol. XXX VIIIIM, No. 4, 687-694.
- Rawat, K. S. and Mukesh Kumar 2009. On Curvature collineations in a Tachibana recurrent space, *Aligarh Bull. Math.*, 28, No. 1-2, 63-69, MR 2769016.
- Rawat, K. S. and Mukesh Kumar 2010. On Weyl-Sasakian bi-recurrent and bi-symmetric spaces, *Jour. PAS*, Vol. 16 (Mathematical Science), PP 484-492.
- Rawat, K. S. and Mukesh Kumar and Nitin Uniyal 2012. On hyperbolically Kaehlerian bi-recurrent and bi-symmetric spaces, *IOSR Jour. of Mathematics (IOSRJM)*, Vol. 1, Issue 2, 28-31. (Included in NASA database).
- Rawat, K. S. and Nitin Uniyal 2011. Lie Derivative of a Linear connexion and various kinds of motions in a Kaehlerian recurrent space of first order, *Jour. of Inter. Aca. of Physical Science*, Vol. 15, No.4., 476-478.
- Rawat, K. S. and Nitin Uniyal 2011. On Conformal transformation in an almost Kaehlerian and Kaehlerian spaces, *Jour. of progressive science*, Vol. 2, No. 2, 138-141.
- Rawat, K. S. and Virendra Prasad 2010. On holomorphically projectively flat parabolically Kaehlerian space, *Rev. Bull. Cal. Math. Soc.*, 18 (1), 21-26.
- Rawat, K.S. and Girish Dobhal 2007. On the bi-recurrent Bochner curvature tensor, *Jour. of the Tensor Society*, Vol. 1, 33-40.
- Rawat, K.S. and Mukesh Kumar 2012. Motions in a Riemannian space and some integrability conditions, *Ultra Scientist of Physical Science (An Inter. Jour. of Physical Science)*, Vol. 24(1) A, 9-14.
- Roter, W. 1974. On Conformally symmetric Ricci-recurrent spaces, *Colloq. Math.* 31, 87-96.
- Singh, S. S. 1971. On Kaehlerian Recurrent and Ricci-recurrent spaces of second order, *Atti Della Accademia della Scienze di Torino*, Vol. 106, PP 509-518.
- Singh, S. S. 1971. On Kaehlerian space with Recurrent Bochner Curvature tensor, *Acc. Naz. Dei. Lincei, Rend, Series VIII*, 51, 3-4, PP 213-220.
- Walker A.G. 1950. On Ruse's Spaces of Recurrent Curvature. *Proc. Land.Math. Soc. II Series* P 52.
- Yano, K. 1965. *Differential Geometry of Complex and almost Complex spaces*, Pergamon press, London.
