



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

ASIAN JOURNAL OF  
SCIENCE AND TECHNOLOGY

Asian Journal of Science and Technology  
Vol. 08, Issue, 10, pp.6272-6277, October, 2017

## REVIEW ARTICLE

### A COMPARATIVE CLINICAL STUDY OF PALASHA KSHARASUTRA AND APAMARGA KSHARASUTRA IN THE MANAGEMENT OF BHAGANDARA (FISTULA-IN-ANO)

Dr. Kunal Kishore, Dr. Munde Bharat Janardan Rao and Dr. Chandrakanth Halli

Dept. Of Shalya Tantra, N.K.J. Ayurvedic Medical College & P.G. Centre, Bidar

#### ARTICLE INFO

##### Article History:

Received 24<sup>th</sup> July, 2017  
Received in revised form  
18<sup>th</sup> August, 2017  
Accepted 20<sup>th</sup> September, 2017  
Published online 30<sup>th</sup> October, 2017

#### ABSTRACT

This research work aims towards study effect of washing on dyed Himalayan Nettle, Cotton and their blends. For this 100% Himalayan Nettle, 100% Cotton and Cotton X Nettle (60 X 40) blends in counts of 3s, 5s, 6s, 7s and 8s are developed. The results obtained are critically analysed.

##### Key words:

Bhagander,  
Fistula in ano.

Copyright©2017, Dr. Kunal Kishore et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

#### INTRODUCTION

Fistula-in-ano is one of the most common ailments pertaining to ano-rectal region. This disease causes discomfort and pain to patient, which creates problems in routine work. In Ayurvedic classics, this disease has been described with the name of BHAGANDARA, which has more similar signs and symptoms with anal fistula and is encountered under Ashtamahagada. As the wound is located in anal region which is more prone to infection, thus takes long time to heal and the condition remains troublesome, operative procedures often leads to complications like recurrences and incontinence.

To alleviate such problems in the management of this disease, it was thought to find out some technique to treat these cases without operative complications. Acharya sushruta in the 17<sup>th</sup> chapter of chikista stana mentioned the application of kshara sutra in Nadi vrana. The Kshara Sutra therapy was practiced and used in since long with great success and without recurrences. The Standard kshara sutra is prepared by repeated coatings of Snuhi ksheera, Apamarga kshara and Haridra. Still some of the problems are faced during the preparation and also in the course of Kshara sutra therapy, like collection and preservation of Snuhi ksheera, burning pain during primary and successive changes. Local irritant skin reactions during course of therapy etc. To overcome these disadvantages was of utmost importance to make the treatment widely popular and acceptable.

In spite of the good rates of cutting, severe pain and burning sensation caused during the treatment with held many patients from accepting this treatment. Thus it gave to many Kshara sutras were tried out. Though each of the thread had good cutting rates and other preparation advantages they also had some disadvantages. Keeping in view the same idea

Palash Ksharasutra<sup>1</sup> was tried in the present study.

#### Aims and Objectives

- Detail and descriptive study of Bhagandara.
- To explore the efficacy of Palash ksharasutra in the management of Bhagandara.

\*Corresponding author: Dr. Kunal Kishore,  
Dept. Of Shalya Tantra, N.K.J. Ayurvedic Medical College & P.G. Centre, Bidar

## MATERIALS AND METHODS

The most important requirement in the clinical study is a well defined protocol. So, in the present study following protocol was followed.

### Source of data

The present clinical study on the management of mutrashmari was carried out at N.K.J.A.M.C. Bidar. This study was carried out at O. P. D. level and the work was limited according to the facilities available in the P. G. Dept. of shalya Tantra .The data was also collected by conducting camps for the purpose of clinical study.

### Selection Criteria

The selection of cases was done on the bases of clinical presentation and the diagnosis was established accordingly. The patients were registered according to the proforma prepared for the study irrespective of their sex, occupation and socio – economic status.

### Inclusive Criteria

- Selection of the patients will be selected 16 -70 age, irrespective sex, religion, occupation, economic status, etc.
- Low anal fistula.

### Exclusion Criteria

- Patients with high rectal fistula.
- Patients suffering from systemic diseases like TB, DM, Ca Rectum, HBsAg, HIV.
- Patients suffering from ulcerative colitis and Crohn's diseases.

### Study Design

Being a clinical study, 30 patients will be selected by Simple Randomized Sampling procedure. All the patients will be screened out by inclusive and exclusive criteria and registered for clinical trial divided in two equal groups.

**Group 1 (Trial Group):** 15 patients of Bhagandara will be treated with Palasha Ksharasutra.

**Group 2 (Control Group):** 15 patients of Bhagandara will be treated with Apamarga Ksharasutra.

In both the groups Ksharasutra will be changed weekly. The duration of the study will be 28 days in total. Patients will be called on every 7<sup>th</sup> days and the cutting rate will be assessed in a specially prepared case sheet. Observations will be analyzed on the basis of assessment parameter (both subjective and objective) critically and scientifically before treatment and after treatment on the 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup> and 28<sup>th</sup> day. Finally the result will be statistically evaluated with the help of paired 't' test within the group for its significance.

### Materials

- Palasha Ksharasutra
- Apamarga Kshara sutra.
- Both the Kshara sutras will be prepared according to the standard method as per Dr. P.J. Deshpande

### Drug Review

Name of the drug	Botanical name	Family	Rasa	Veerya	Vipaka	Guna	Karma
Palasha	Butea Monospermia	Leguminosae	Katu Tikta kashaya	Ushna	Madhura	Ruksha laghu	Kapha-Vata shamaka,
Snuhi	Euphorbia nerifolia	Euphorbiaceae	Katu	Ushna	Katu	Laghu- teekshna	Kapha- vata hara, Deepana, rechana
Haridra	Curcuma longa	Scitamineaceae	Tikta, katu	Ushna	Katu	Ruksha, laghu	Kapha vatahara, lekhana, vishagna, varnya.
Apamarga	Achyranthus aspera linn	Amaranthaceae	Katu Tikta	Ushna	Katu	Laghu Ruksha Tikshna Sara	Shoothara , Vedana sthapana, Vishaghna, Vrana Shadhana, Twaga doshahara.

### Assesment Criteria

**Subjective criteria:** Pain according to

- Grade 0 (-) -Absolute No pain  
 Grade 1 (+) -Mild pain  
 Grade 2 (++) -Moderate pain  
 Grade 3 (+++) -Severe pain

**Itching**

Grade 0 (-)	-No itching
Grade 1 (+)	-Itching sometimes but not interfering with activities can be controlled voluntarily
Grade 2 (++)	-Itching interference with function involuntary and uncontrolled associated with skin patches.
Grade 3 (+++)	-Itching that disturbs the sleep and or demand treatment associated with skin patches.

**Discharge**

Grade 0 (-)	-	No Discharge
Grade 1 (+)	-	Mild discharge single pad is sufficient per day.
Grade 2 (++)	-	Moderate discharge 2-3 pads necessary per day
Grade 3 (+++)	-	Profuse discharge more than 3 pads are necessary per day

**Objective Criteria**

**Local tenderness:** was assessed by palpation of anal region.

G<sub>0</sub> – No tenderness

G<sub>1</sub> – mild – tender but can be palpated.

G<sub>2</sub>- moderate – tender on gentle palpation.

G<sub>3</sub>- Severe – patient denies touching.

**Induration**

G<sub>0</sub> - Absent

G<sub>1</sub> - Slight swelling around the wound margin without indurations.

G<sub>2</sub> - Swelling around wound margin with little area of indurations.

G<sub>3</sub> - Swelling with marked indurations.

**Length of tract (in cm)**

**Unit cutting time:** Initial length of the tract – length of the tract remaining

No of weeks taken.

**Investigation Required**

Blood Routine & Urine Examination.

Culture and Sensitivity of wound discharge (if necessary),

Fistulography (if necessary),

**RESULTS****Trial group result-1**

Sign and Symptoms	BT	Mean ± S.E.	Assesment	Mean ± S.E.	Df	T-value	p-value	p-value	Effectiveness %	Remark					
Pain on VAS	2.06±0.18		1	2.06±0.18	14	0.56	0.58	>0.05	2.85	NS					
			2	1.86±0.13							3.68	0.0003	<0.001	27.77	HS
			3	1.53±0.16							15.58	3.16E	<0.001	27.77	HS
			4	0.86±0.13							18.66	1.32E-11	<0.001	48.57	HS
Swelling	2.21±0.18		1	2.21±0.18		1.46	0.16	>0.05	6.66	NS					
			2	2.21±0.18							1.46	0.16	>0.05	6.66	NS
			3	1.21±0.10							6.51	1.37E-05	<0.001	45.16	HS
			4	0.42±0.13							14.50	7.91-10	<0.001	80.64	HS
Itching	2.06±0.18		1	2.06±0.18		0.56	0.58	>0.05	3.12	NS					
			2	2.06±0.18							0.56	0.58	>0.05	3.12	NS
			3	1±0.13							9.02	28E-07	<0.001	51.61	HS
			4	0.4±0.13							10.45	35E-08	<0.001	80.64	HS
Discharge	2.33±0.18		1	2.33±0.18		2.25	0.040	<0.05	10.52	S					
			2	1.73±0.15							4.58	0.0004	<0.001	25.714	HS
			3	1.2±0.10							6.85	7.82E-06	<0.001	48.57	HS
			4	0.46±0.13							20.54	7.46-12	<0.001	80	HS
Local Tenderness	2.4±0.13		1	2.4±0.13		1	0.066	>0.05	2.85	NS					
			2	1.73±0.15							5.29	0.00014	<0.001	27.77	HS
			3	1.2±0.10							11.22	2.19E-08	<0.001	50	HS
			4	0.66±0.12							14.66	6.85E-6	<0.001	72.22	HS
Length of Track	3.46±0.17		1	3.46±0.17		10.66	1.53E-07	<0.001	28.03	HS					
			2	2.50±0.17							10.26	6.72E-08	<0.001	25.8	HS
			3	1.57±0.14							17.80	5.16E	<0.001	54.52	HS
			4	0.57±0.10							19.55	1.46E-11	<0.001	83.42	HS

The above statistical analysis shows that in case of

**Pain:** The mean  $\pm$  S.E. before treatment was  $2.06 \pm 0.18$  and was same  $2.06 \pm 0.18$  after 7 days, was reduced to  $1.86 \pm 0.13$  after 14 days, and  $1.53 \pm 0.16$  after 21 days.  $0.86 \pm 0.13$  after 28 days. The test of significance shows that palash is highly Significant to reduce pain with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

**Swelling:** The mean  $\pm$  S.E. before treatment was  $2.21 \pm 0.18$  and was same  $2.21 \pm 0.18$  after 7 days, was reduced to  $2.21 \pm 0.18$  after 14 days, and  $1.21 \pm 0.10$  after 21 days.  $0.42 \pm 0.13$  after 28 days. The test of significance shows that palash is highly Significant to reduce swelling with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

**Itching:** The mean  $\pm$  S.E. before treatment was  $2.06 \pm 0.18$  and was same  $2.06 \pm 0.18$  after 7 days, was same to  $2.06 \pm 0.18$  after 14 days, and was reduced to  $1 \pm 0.13$  after 21 days.  $0.4 \pm 0.13$  after 28 days. The test of significance shows that palash is highly Significant to reduce itching with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

**Discharge:** The mean  $\pm$  S.E. before treatment was,  $2.33 \pm 0.18$  and was same  $2.33 \pm 0.18$  after 7 days, was reduced to  $1.73 \pm 0.15$  after 14 days, and  $1.2 \pm 0.10$  after 21 days.  $0.46 \pm 0.13$  after 28 days. The test of significance shows that palash is highly Significant to reduce Discharge with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

**Local of Tenderness:** The mean  $\pm$  S.E. before treatment was,  $2.4 \pm 0.13$  and was same  $2.4 \pm 0.13$  after 7 days, was reduced to  $1.73 \pm 0.15$  after 14 days, and  $1.2 \pm 0.10$  after 21 days.  $0.66 \pm 0.12$  after 28 days. The test of significance shows that palash is highly Significant to reduce Local tenderness with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

**Length of track:** The mean  $\pm$  S.E. before treatment was,  $3.46 \pm 0.17$  and was same  $3.46 \pm 0.17$  after 7 days, was reduced to  $2.50 \pm 0.17$  after 14 days, and  $1.57 \pm 0.14$  after 21 days.  $0.57 \pm 0.10$  after 28 days. The test of significance shows that palash is highly Significant to reduce Length of track with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

#### Effectiveness % of Trial Group-1

##### Control group result: 2

Sign and Symptoms	BT Mean $\pm$ S.E.	Assesment	Mean $\pm$ S.E.	Df	T-value	p-value	p-value	Effectiveness %	Remark
Pain on VAS	1.53 $\pm$ 0.16	1	1.53 $\pm$ 0.16	14	4.67	0.66	>0.05	2.85	NS
		2	1.73 $\pm$ 0.15		2.25	0.002	>0.001	10.52	S
		3	1.26 $\pm$ 0.15		7.29	0.00011	<0.001	36.84	HS
		4	0.73 $\pm$ 0.11		8.76	0.000131	<0.001	52.94	HS
Swelling	2.13 $\pm$ 0.16	1	2.13 $\pm$ 0.16		1.46	0.16	>0.05	6.06	NS
		2	2.06 $\pm$ 0.18		0.56	0.58	>0.05	3.12	NS
		3	1.13 $\pm$ 0.135		7.24	4.25E-06	<0.001	46.8	HS
		4	0.2 $\pm$ 0.10		16.35	1.61E-10	<0.001	90.625	HS
Itching	2.06 $\pm$ 0.18	1	2.06 $\pm$ 0.18		1.87	0.08	>0.05	9.67	NS
		2	1.86 $\pm$ 0.13		1.87	0.082	>0.05	9.67	NS
		3	0.86 $\pm$ 0.13		8.29	9.02E-07	<0.005	58.06	S
		4	0.2 $\pm$ 0.10		9.72	1.13E-07	<0.001	90.32	HS
Discharge	2.13 $\pm$ 0.13	1	2.13 $\pm$ 0.13		0.56	0.58	>0.05	3.03	NS
		2	1.6 $\pm$ 0.13		4	0.0013	>0.001	25	S
		3	1 $\pm$ 0		8.5	6.71E-07	<0.001	53.12	HS
		4	0.2 $\pm$ 0.10		12.61	4.92E-09	<0.001	90.62	HS
Local Tenderness	2.4 $\pm$ 0.13	1	2.4 $\pm$ 0.13		1	0.33	>0.05	2.85	NS
		2	1.6 $\pm$ 0.13		5.24	0.000124	<0.001	35.13	HS
		3	1.33 $\pm$ 0.09		10.58	4.61E-08	<0.001	54.05	HS
		4	0.33 $\pm$ 0.12		23.48	1.21	<0.001	86.48	HS
Length of Track	4.2 $\pm$ 0.25	1	4.2 $\pm$ 0.25		2.61	0.02	<0.05	21.18	S
		2	3.10 $\pm$ 0.25		9.66	1.43E-07	<0.001	26.03	HS
		3	1.97 $\pm$ 0.25		1.67	1.17E-10	<0.001	53.01	HS
		4	0.82 $\pm$ 0.19		29.56	5.09E-14	<0.001	80.47	HS

The above statistical analysis shows that in case of

**Pain:** The mean  $\pm$  S.E. 1.53 $\pm$ 0.16 before treatment was same 1.53 $\pm$ 0.16 after 7 days, was reduced to 1.73 $\pm$ 0.15 after 14 days, 1.26 $\pm$ 0.15 and after 21 days. 0.73 $\pm$ 0.11 after 28 days. The test of significance shows that Apamarga is highly Significant to reduce pain with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

**Swelling:** The mean  $\pm$  S.E. before treatment was 2.13 $\pm$ 0.16 and was same 2.13 $\pm$ 0.16 after 7 days, was reduced to 2.06 $\pm$ 0.18 after 14 days, and 1.13 $\pm$ 0.135 after 21 days. 0.2 $\pm$ 0.10 after 28 days. The test of significance shows that Apamarga highly Significant to reduce swelling with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

**Itching:** The mean  $\pm$  S.E. before treatment was 2.06 $\pm$ 0.18 and was same 2.06 $\pm$ 0.18 after 7 days, was reduced to 1.86 $\pm$ 0.13 after 14 days, and 0.86 $\pm$ 0.13 after 21 days. 0.2 $\pm$ 0.10 after 28 days. The test of significance shows that Apamarga is highly Significant to reduce itching with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

**Discharge:** The mean  $\pm$  S.E. before treatment was,  $2.13 \pm 0.13$  and was same  $2.13 \pm 0.13$  after 7 days, was reduced  $1.6 \pm 0.13$  after 14 days, and  $1 \pm 0$  after 21 days.  $0.2 \pm 0.10$  after 28 days. The test of significance shows that Apamarga is highly Significant to reduce Discharge with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

**Local of Tenderness:** The mean  $\pm$  S.E. before treatment was,  $2.4 \pm 0.13$  and was same  $2.4 \pm 0.13$  after 7 days, was reduced to  $1.6 \pm 0.13$  after 14 days, and  $1.33 \pm 0.09$  after 21 days.  $0.33 \pm 0.12$  after 28 days. The test of significance shows that Apamarga is highly Significant to reduce Local tenderness with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

**Length of track:** The mean  $\pm$  S.E. before treatment was,  $4.2 \pm 0.25$  and was same  $4.2 \pm 0.25$  after 7 days, was reduced  $3.10 \pm 0.25$  after 14 days, and  $1.97 \pm 0.25$  after 21 days.  $0.82 \pm 0.19$  after 28 days. The test of significance shows that Apamarga is highly Significant to reduce Length of track with the P-value  $< 0.001$  in AT1, AT2, AT3&AT4 respectively.

## Effectiveness % of Control Group-2

### Overall clinical assessment of therapy

In group I, 20 patients (80%) complete cured. In group II, 20 patients (90%) got complete cured Finally overall assessment of the therapy was analyzed clinically, Where at the 1<sup>st</sup> follow up on 7<sup>th</sup> day in group - I Moderate improvement in 5 patients i.e. (25%). Mild improve in 6 patients i.e. (30%) and not improvement was noticed in 9 patients i.e. (45%) where as group- II Moderate improvement in 6 patients i.e. (30%). Mild improve in 5 patients i.e. (25%) and not improvement was noticed in 9 patients i.e. (45%), Where at the 2<sup>st</sup> follow up on 14<sup>th</sup> day in group - I Max improvement in 2 patients i.e. (10%). Moderate improvement in 3 patients i.e. (15%). Mild improve in 4 patients i.e. (20%) and not improvement was noticed in 6 patients i.e. (30%) where as group- II Max improvement in 0 Patients i.e. (0%). Moderate improvement in 5 patients i.e. (25%). Mild improve in 6 patients i.e. (30%) and not improvement was noticed in 3 patients i.e. (15%) Where at the 3<sup>rd</sup> follow up on 21<sup>st</sup> day in group - I 1 patients i.e. (5%) was Cured and Max improve in 5 patients i.e. (25%). Moderate improvement in 7 patients i.e. (35%). Mild improve in 5 patients i.e. (25%) and not improvement was noticed in 2 patients i.e. (10%). Where as group - II 4 patients i.e. (20%) was Cured and Max improve in 8 patients i.e. (40%). Moderate improvement in 3 patients i.e. (15%) Mild improve in 4 patients i.e. (20%) and not improvement was noticed in 1 patients i.e. (15%).

Whereas the 4<sup>th</sup> follow up on 28<sup>st</sup> day in group - I 16 patients i.e. (80%) was Cured and Max improve in 4 patients i.e. (20%). group - II 18 patients i.e. (90%) was Cured and Max improve in 2 patients i.e. (10%). After comparison of all above figures of sign & symptoms in Trial and Control groups, we can conclude that the treatment used in control group i.e. Treatment of Bhagandara (Fistula-in-ano) by Apamarga ksharasutra is more effective as compared to the treatment trail group Arka ksharasutra.

### Over all clinical assessment of therapy

	GROUP-I A.T.								GROUP-II A.T.							
	7th day		14th day		21st day		28th day		7th day		14th day		21st day		28th day	
	No. of pt	%	No. of pt	%	No. of pt	%	No. of pt	%	No. of pt	%	No. of pt	%	No. of pt	%	No. of pt	%
Cured -100%	0	0%	0	0%	1	5%	16	80%	0	0%	0	0.00%	4	20%	18	90%
Max. Improved (75% to 99%)	0	0%	2	10%	5	25%	4	20%	0	0%	5	25%	8	40%	2	10%
Mod. Improved (50% to 74%)	5	25%	3	15%	7	35%	0	0%	6	30%	6	30%	3	15%	0	0%
Mild Improved (25% to 49%)	6	30%	4	20%	5	25%	0	0%	5	25%	6	30%	4	20%	0	0%
Not Improved (<25%)	9	45%	6	30%	2	10%	0	0%	9	45%	3	15%	1	5%	0	0%

## DISCUSSION

After comparison of all above figures of signs & symptoms in Trial and Control groups, we can conclude that the treatment used in control group i.e. Treatment of Bhagandara (Fistula-in-ano) by Apamarga ksharasutra is more effective as compared to the treatment with trail group i.e. Palashksharasutra.

### Conclusions

After a clinical observation and statistical evaluation the following conclusions were drawn.

- The management of fistula in ano by khara sutra has been proved effective by this study.
- The undersired effects of ksharasutra management could be minimized by using palash ksharasutra..
- Palash kshara sutra has been found very effective in relieving symptoms i.e pain, swelling, itching, discharge ,and local tenderness in fistula in shorter time.
- Both the varieties are cost effective can be easily prepared and can be easily applied with less recurrence after treatment.
- Management with Apamarga ksharasutra shows superiority than Palash ksharasutra.

**REFERENCES**

- Indra Tripathi, Dr. and Dayashankar Tripathi, Dr. Yogarathnakar with Vidyapradha Hindi Commentary, Pub. by Krishnadas Academy, Varanasi, 1<sup>st</sup> Ed. 1998.
- Madhava Nidana with Madhukosha Sanskrit Commentery vol-2, Chaukhamba Sanskrit Bavan- Varanasi, Ma. Ni.-46/1 page no.-134 .
- Shastri Kaviraj Ambika Dutta, Sushruta samhita, Chikitsa Sthan, Choukhamba sansthan publishers, Varanasi, Edition 2010, Su. Ni.- 4/3 page no.-316.
- Shastri Kaviraj Ambika Dutta, Sushruta samhita, Chikitsa Sthan, Choukhamba sansthan publishers, Varanasi, Edition 2010, Su. Chi.- 17/13 page no.101.
- Shastri Kaviraj Ambika Dutta, Sushruta samhita, Chikitsa Sthan, Choukhamba sansthan publishers, Varanasi, Edited reprint 2006, 8:45<sub>pp</sub>
- Shastri Kaviraj Ambika Dutta, Sushruta samhita, Niodan Sthan, Choukhamba sansthan publishers, Varanasi, Edition 2010, Su. Ni.4/4, page no. – 317.
- Shastri Kaviraj Ambika Dutta, Sushruta samhita, Sutra Sthan, Choukhamba sansthan publishers, Varanasi, Edition 2006, Su. Su.11/12, page no.-47.

\*\*\*\*\*