



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

ASIAN JOURNAL OF
SCIENCE AND TECHNOLOGY

Asian Journal of Science and Technology
Vol. 11, Issue, 09, pp.11182-11186, September, 2020

RESEARCH ARTICLE

CHANGES IN CHEMICAL COMPOSITION AND FUNCTIONAL PROPERTIES OF A FORTIFIED IVORIAN LOCAL FOOD BASSI OF MILLET

Niaba Koffi Pierre Valery^{1,3*}, COMBO Agnan Marie-Michel³, Deffan Zranseu Ange Benedicte³,
Marina C. Kouakou⁴, Germain A. Brou² and Gildas K. Gbassi^{1,2}

¹Food, Water and Beverage Control Service, National Laboratory of Public Health (LNSP), BP 2403 Abidjan 18,
Côte d'Ivoire

²Department of Analytical Chemistry, General and Mineral Chemistry, UFR Pharmaceutical and Biological
Sciences, Félix Houphouët Boigny University, BPV 34 Abidjan 01, Cote d'Ivoire

³Agro-valorization Laboratory, Agro forestry Department, Jean Lorougnon Guédé University,
Daloa, Côte d'Ivoire, BP 150

⁴Nutrition and Food Security Laboratory, Nagui Abrogoua University, Abidjan, Côte d'Ivoire

ARTICLE INFO

Article History:

Received 17th June, 2020

Received in revised form

09th July, 2020

Accepted 24th August, 2020

Published online 30th September, 2020

ABSTRACT

Six (6) formulations of "BASSI" based on a mixture of millet and soybeans have been developed. Soy flour was incorporated in amounts of 0, 10, 20, 30, 40 and 50%. The results showed that the nutritional value significantly increases with the incorporated soy rate, except the carbohydrate content which decreases. The water absorption capacity, oil absorption capacity and emulsifying activity also increased. As for the emulsion stability and the bulk density, the values decrease. The incorporation of soy improves the nutritional and functional properties of "BASSI".

Key words:

"BASSI", Cereal-Based Food, Chemical
Composition, Functional Properties.

Citation: Niaba Koffi Pierre Valery, COMBO Agnan Marie-Michel *et al.* 2020. "Changes in chemical composition and functional properties of a fortified ivoirian local food "bassi" of millet", *Asian Journal of Science and Technology*, 11, (09), 11182-11186.

Copyright © 2020, Niaba Koffi Pierre Valery *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Dietary deficiencies are major scourges news. According to FAO, more than one billion people are affected by malnutrition in worldwide, most of which is in developing countries (FAO, 2009). The main causes are a protein-energy food deficit and a deficiency of certain key micronutrients such as calcium, iron and zinc (Soro *et al.*, 2013). But we should add the unsustainable and inequitable access to food by the most vulnerable social strata and also the bad nutritional practices. Grains, including maize and millet are the staple food of many populations around the world. In Cote d'Ivoire, maize is the second cereal most cultivated and consumed after rice (Akanvou *et al.*, 2006), while millet is the 3rd (Békoye and Akanvou, 2005). These food crops are used in the manufacture of many traditional dishes, including "BASSI".

This multicultural food made from local product is consumed by the Ivorian population and several countries of the sub region. It is obtained after cooking cornmeal or millet and added sugar. But its nutritional value seems to be unbalanced because the main raw materials (millet) for the manufacture of this commodity are rich in carbohydrates and low in proteins. The levels are 73% and 9.2% for maize and 67% and 11.8% for millet respectively (FAO, 1995). Generally, grains have low protein concentration and the quality of these proteins is limited by deficiencies in certain essential amino acids. Thus, it is necessary to incorporate a product with high nutritional density, and soy appears to meet this requirement. In most developing countries, soybean cultivation is encouraged. This legume is known for its seeds rich in protein (40%) and fat (20%), giving him the rank of first oleaginous-protein cultivated in the world (Duc *et al.*, 2010). According to Zannou-Tchoko *et al.* (2011), soybean contains in balanced proportions proteins of good biological value with all essential amino acids. It also contains fatty acids such as omega (Demaison and Moreau, 2002). In addition to provide essential nutrients, it is used for its excellent functional properties in

***Corresponding author:** Niaba Koffi Pierre Valery,
Food, Water and Beverage Control Service, National Laboratory of
Public Health (LNSP), BP 2403 Abidjan 18, Côte d'Ivoire.

improving food processes (Wolf, 1970). Several formulations of "BASSI" from millet with soybeans incorporating were developed with the objective to determine the chemical composition of such mixtures; and assess their functional properties.

MATERIALS AND METHODS

Materials

The plant material used was pearl millet (*Pennisetum glaucum*) and yellow soybeans (*Glycine max*).

Methods

Formulations of "BASSI" improved with soybean: Twelve formulations of "BASSI" containing soybean have been developed, six from millet. Soy flour was mixed with that of millet (IF) in the following proportions: 0% (IF0), 10% (IF1), 20% (IF2), 30% (IF3), 40% (IF4) and 50% (IF5). 25 g of sugar were added to 100 g of each mixture of flour. A specific volume of water was added to each mixture. The preparation process of "BASSI" is presented in figure 1.




Figure 1: Scheme of "BASSI" manufacturing improved with soybean

Chemical analysis of "BASSI" improved with soybean:

Water content and ashes content were determined according to an official method (AOAC, 1990). Moisture was determined by drying in an oven at 105 °C for 24 hours and ash using a muffle furnace at 550 °C for 24 h. Proteins were quantified according to AOAC (1990) using the Kjeldahl method which taking place in three steps namely mineralization, distillation and acid-base quantification. Fat content was determined according to the method of Folch *et al.* (1957) using vacuum filtration and a solvent combination (chloroform and methanol) to extract the fat. Total carbohydrate was determined by difference of the total material to other biochemical compounds. Energy value corresponding to the available energy was calculated using the specific coefficients of Atwater and Benedict (1902) for proteins, lipids and carbohydrates. The pH and total acidity were determined according to AOAC (1990). For the last two parameters, each sample was diluted in 100 ml of distilled water, stirred for 30 min and the resulting mixture was filtered before analysis.

Functional properties of "BASSI": The water absorption capacity (WAC) was determined using the method of Phillips *et al.* (1988). The oil absorption capacity (OAC) was determined according to the method of Sosulski (1962). The hydrophilic-lipophilic index (HLI) as defined by Njintang *et al.* (2001) was calculated as the ratio of WAC/OAC. The emulsifying activity (EA) and emulsion stability (ES) were determined by the method of Yasumatsu *et al.* (1972). The bulk density (BD) was determined according to the method of Narayana and Narasimha (1982).

Statistical data processing: All measurements were performed in triplicate. Statistical differences were verified by ANOVA with SPSS 19.0 software. The comparison of means was carried out with the Duncan test for a significance level set at 5%.

RESULTS AND DISCUSSION


Results

Chemical analysis of "BASSI" improved with soybean

Case of "BASSI" based millet: The nutrient composition of all formulations is presented in table 1. Water content is between 4.34 ± 0.01 and 4.70 ± 0.02 g/100 g of DM. Protein, fat and ash increases with incorporated soy rate ($p < 0.05$). The mineral composition gradually increases with the incorporation of soybean. The macro-minerals and trace elements increase significantly ($p < 0.05$). The total carbohydrate content decreases ($p < 0.05$). The incorporation of soybean improves the energy value of formulations ($p < 0.05$). The results concerning sugar content, pH and acidity are shown in table 2. Total sugar content decreases as soybean increases. Values range between 15.73 ± 0.56 and 14.36 ± 0.46 mg/ml. pH and acidity changing inversely as soybean increases. The pH values are significantly different ($p < 0.05$). Acidity is statistically identical for all formulation.

Functional properties of "BASSI" improved with soybean

Case of "BASSI" based millet: The results for functional properties are shown in Table 3.



Photocopy: "BASSI" improved with soybean

Table 1. Nutritional value of "BASSI"based millet improved with soybean

Parameters	Content (g/100 g of dry matter)					
	IF0	IF1	IF2	IF3	IF4	IF5
Moisture	4.68 ^c ± 0.03	4.60 ^d ± 0.01	4.34 ^a ± 0.01	4.38 ^b ± 0.10	4.48 ^e ± 0.02	4.70 ^e ± 0.02
Proteins	7.10 ^a ± 0.02	10.54 ^b ± 0.39	14.42 ^c ± 0.02	17.45 ^d ± 0.03	20.74 ^e ± 0.03	24.78 ^f ± 0.02
Fat	4.75 ^a ± 0.04	5.41 ^b ± 0.07	6.27 ^c ± 0.13	7.90 ^d ± 0.08	9.74 ^e ± 0.22	10.42 ^f ± 0.05
Ashes	1.11 ^a ± 0.01	1.27 ^b ± 0.00	1.87 ^c ± 0.01	2.09 ^d ± 0.10	2.63 ^e ± 0.00	2.83 ^f ± 0.01
Carbohydrates	82.39 ^f ± 0.14	78.19 ^e ± 0.38	73.11 ^d ± 0.13	68.18 ^c ± 0.06	62.41 ^b ± 0.19	57.28 ^a ± 0.07
Energetic value*	400.74 ^a ± 0.24	403.61 ^b ± 0.32	406.52 ^c ± 0.66	413.65 ^d ± 0.41	420.23 ^e ± 1.20	422.11 ^f ± 0.50
Minerals	Content (mg/100 g of dry matter)					
Calcium	28.28 ^a ± 0.16	32.75 ^b ± 0.20	34.54 ^c ± 0.01	40.29 ^d ± 0.18	46.11 ^e ± 0.20	53.08 ^f ± 0.07
Phosphorus	89.47 ^a ± 0.51	93.75 ^b ± 0.12	100.54 ^c ± 0.13	110.79 ^d ± 0.44	131.83 ^e ± 0.71	157.44 ^f ± 0.09
Potassium	1091.00 ^a ± 2.71	1136.10 ^b ± 4.03	1185.93 ^c ± 1.97	1191.72 ^c ± 9.32	1206.01 ^d ± 3.01	1309.02 ^e ± 7.74
Magnesium	69.08 ^a ± 0.28	79.48 ^b ± 1.15	83.38 ^c ± 0.95	99.63 ^d ± 2.11	125.11 ^e ± 0.77	137.64 ^f ± 0.12
Sodium	71.96 ^a ± 0.22	83.42 ^b ± 0.31	89.72 ^c ± 0.10	104.47 ^d ± 0.30	106.34 ^e ± 0.34	126.80 ^f ± 0.19
Manganese	0.39 ^a ± 0.02	0.55 ^b ± 0.04	0.67 ^c ± 0.03	0.68 ^c ± 0.00	0.79 ^d ± 0.01	1.11 ^e ± 0.00
Iron	4.65 ^a ± 0.06	4.78 ^b ± 0.03	4.86 ^b ± 0.03	5.85 ^c ± 0.03	5.90 ^c ± 0.00	6.12 ^d ± 0.05
Copper	0	0	0	0	0	0
Zinc	0.97 ^a ± 0.01	1.04 ^a ± 0.12	1.05 ^a ± 0.02	1.26 ^b ± 0.02	1.28 ^b ± 0.05	1.61 ^c ± 0.03

Table 2. Total sugars, pH and acidity of "BASSI" based millet improved with soybean

Parameters	IF0	IF1	IF2	IF3	IF4	IF5
Total sugars*	15.73 ^c ± 0.56	15.38 ^c ± 0.29	15.31 ^c ± 0.12	15.10 ^{bc} ± 0.20	14.50 ^{ab} ± 0.19	14.36 ^a ± 0.46
pH	5.43 ^a ± 0.00	6.13 ^b ± 0.00	6.21 ^c ± 0.38	6.37 ^d ± 0.00	6.47 ^e ± 0.00	6.50 ^f ± 0.00
Acidity**	3.33 ^a ± 0.57	2.67 ^a ± 0.57	2.67 ^a ± 0.57	2.33 ^a ± 0.57	2.33 ^a ± 0.57	2.33 ^a ± 0.57

Values followed by different letters in a line are statistically different ($\alpha = 0.05$). Values represent the mean \pm SD of three independent measurements ($n = 3$) * (mg/ml) ** (meq-g/100 g of dry matter).

Table 3. Functional properties of "BASSI" based millet improved with soybean

Parameters	IF0	IF1	IF2	IF3	IF4	IF5
WAC*	269.86 ^a ± 2.08	278.50 ^b ± 1.46	287.30 ^c ± 0.89	292.67 ^d ± 0.29	294.43 ^{dc} ± 0.25	295.44 ^e ± 0.27
OAC**	158.42 ^a ± 0.47	161.63 ^b ± 0.71	166.30 ^c ± 0.42	167.77 ^d ± 0.61	169.37 ^e ± 1.28	172.75 ^f ± 0.56
HLI	1.70 ^a ± 0.01	1.72 ^{bc} ± 0.00	1.42 ^c ± 0.01	1.73 ^c ± 0.01	1.74 ^c ± 0.00	1.71 ^{ab} ± 0.00
EA (%)	59.50 ^a ± 0.12	61.55 ^b ± 1.27	65.04 ^c ± 1.61	68.94 ^d ± 0.10	71.52 ^e ± 1.17	76.02 ^f ± 0.74
ES (%)	96.66 ^c ± 1.15	95.35 ^d ± 0.10	93.38 ^e ± 0.14	93.29 ^e ± 0.00	69.13 ^b ± 1.77	35.00 ^a ± 1.44
BD (g/ml)	0.81 ^f ± 0.00	0.78 ^e ± 0.00	0.77 ^d ± 0.01	0.75 ^c ± 0.01	0.71 ^b ± 0.01	0.66 ^a ± 0.01

Values followed by different letters in a line are statistically different ($\alpha = 0.05$). WAC= Water Absorption Capacity, OAC = Oil Absorption Capacity, HLI = Hydrophilic Lipophilic Index, EA = Emulsifying Activity, ES = Emulsion Stability, BD = Bulk Density * (g water/100 g of dry matter), ** (g oil/100 g of dry matter).

The WAC, OAC and EA increase with soybean incorporation. Values are significantly different ($p < 0.05$). As for the ES, the values decrease and vary from 96.66 ± 1.15 to $35.00 \pm 1.44\%$. The values of HLI are between 1.42 ± 0.01 and 1.74 ± 0.00 and are significantly identical for the formulations at 30% (IF3) and 40% (IF4). The BD decreases gradually as soybean increases and values are significantly different.

DISCUSSION

Chemical analysis of "BASSI" improved with soybean: The protein content of the formulations increases with soybean incorporation. This result is in agreement with those of Akubor and Onimawo (2003) and Akusu and Wordu (2013) in their studies on the functional properties of local products. The protein content (10.54%) of IF1 formulation is lower than that found by Kalimbira *et al.* (2004) which obtained 14.15%. This difference could be related to the millet variety used. According to Sanogo (1994) *Codex Alimentarius* recommends a minimum of 13% protein for infant foods. IF2, IF3 and IF5 formulations are in compliance with Codex recommendations. The fat content also increases with soybean incorporation because soybean is rich in fat. Millet flours are naturally low in fat (4.75% for millet). Incorporation of soybean which is an oleaginous improves the fat content of the food. According to Demaison and Moreau (2002), the omega fats present in soybean are responsible for cardiovascular and immune balance. IF3, IF4 and IF5 formulations exceed the recommendations of the *Codex Alimentarius* (7%) for baby food (Sanogo, 1994). The ash content reflects the mineral composition of a food. That of our formulations increases with soybean rate. The formulations (IF3, IF4, IF5) exceed the recommendations of *Codex Alimentarius* (2%) for baby food (Sanogo, 1994). The results presented in tables 1 and 2 show that with the exception of calcium and copper (only for formulations containing millet), the mineral content of the formulations from 10 to 50% tend to satisfy the minimum desirable contents of an infantile flour (Mouquet *et al.*, 1988).

All formulations are excellent sources of carbohydrates. Formulations of 0 to 30% of soybean for both millet exceed the recommendations of the *Codex Alimentarius* (68%) for baby food (Sanogo, 1994). The energy value increases as soybean increases. This could be attributed to high levels of fat content (Kalimbira *et al.*, 2004). All formulations have energy values similar to the recommendations of *Codex Alimentarius* (400 Kcal/100 g of DM) for baby food (Sanogo, 1994). The pH values are statistically different while those of acidity are identical. Millet-based formulations are acidic. These acidic pH values are beneficial for food, especially in infant food (Soro *et al.*, 2013) as most enzyme reactions are favored in acidic media.

Functional properties of "BASSI" improved with soybean

The oil absorption capacity (WAC) provides information on the maximum amount of water absorbed by food (Siddiq *et al.*, 2010). WAC of formulations increases with soybean rate. These results are consistent with the work of Akubor and Badifu (2004) which showed that increasing the WAC could be explained by the increase in protein content (many hydrophilic residues). WAC values for the formulations from 20 to 50% (287.30 to 295.44% for millet) are superior to those obtained by Akubor and Onimawo (2003) (180-190 %). High values of WAC can also be assigned to cooking effect.

Cooking would lead to dissociation and modification of protein molecules in monomeric subunits that can retain more water in the binding sites as suggested by Lin *et al.* (1974). A high value of WAC of flour is an important property in pastry flour (Wolf, 1970).

The oil absorption capacity (OAC) is important for the development of flavor and food preservation, because it would prevent the development of oxidative rancidity in reducing the availability of oil for oxygen in food (Siddiq *et al.*, 2010). In our study, the OAC increases with soybean incorporation. This increase is attributed to their protein content and the cooking effect as suggested by an author (El-Adawy, 2000). The type of protein can also influence oil retention properties, because the oil is attached to the nonpolar side chains of proteins as mentioned by some authors (Siddiq *et al.*, 2010) in their work on the physical and functional characteristics of beans flour. The ability of proteins to retain oil is an interesting property because it allows good retention of flavor in the food processes, improving palatability (Moure *et al.*, 2006). The flours of this study could be good lipophilic constituents and therefore suitable for preparing soups and cakes as shown by some authors (Aremu *et al.*, 2007).

The hydrophilic-lipophilic index (HLI) assesses the comparative flour affinity for water and oil. The values for the formulations containing millet (1.42 and 1.74) are superior to those reported by Njintang *et al.* (2001) for cowpea (about 1.12). More HLI approach to value 1, more the flour has good affinity for water as well as oil. All the values are greater than 1, this means that the flours have more affinity for water than oil. The emulsifying activity (EA) is an important parameter in the development of formulations requiring mixtures of water and oil, such as baked goods (Shad *et al.*, 2013). This parameter increases with soy incorporation. Improving the EA could be attributed to the increase in protein content as suggested by some authors (Moure *et al.*, 2006). Soy protein helps the formation of emulsions and their stability during food processing. Our results agree with those of Siddiq *et al.* (2010) which reported in their study that the high stability of flour emulsion was due to the globular nature of the main soy proteins. The bulk density (BD) is influenced by the size of the starch particles. The resulting values decrease as soybean increases. This could probably be due to the fact that millet have higher densities than soybeans (Akubor and Onimawo, 2003). The same authors reported a similarity with the cookies made from soy flour and maize. This decrease may be also due to the small amount of starch in soya meal compared to millet. This property is important during mixing and packaging (Sakai, 1979).

Conclusion and perspectives

This study is a contribution for improving the nutritional quality and functional properties of "BASSI", a local product based on a mixture of millet and soybean. "BASSI" enrichment with soybean improves nutrient levels and functional properties of the formulations. "BASSI" consumption can reduce the risk of malnutrition among the population, particularly in children. The results for the functional properties provide useful data for their potential uses in food industry. In perspective, it would be interesting to determine the profiles of amino acids and fatty acids that are crucial for assessing the quality of protein and fat. In addition, the

establishment of an extension policy is needed to bring the population to consume improved traditional dishes that meet the nutritional quality requirements.

Acknowledgements

The technical support of Laboratoire National de la Santé Publique (LNSP) is acknowledged. The author wishes to express sincere appreciation to Prof. Malan Kla Anglade for its technical comments and manuscript editing.

REFERENCES

- Agbo NG. 1996. Supplementation of a traditional Ivoirian food (Attieke) with soybean. These de doctorat: Université de Cocody-Abidjan, p 15-20.
- Akanvou L, Akanvou R, Anguété K, Diarrassouba L. 2006. Bien cultiver le maïs en Côte d'Ivoire. Centre National de Recherche Agronomique (CNRA); 4 p.
- Akubor PI, Onimawo, IA. 2003. Functional properties and performance of soybean and maize flour blends in cookies. *Plant Foods Hum. Nutr.* 58:1-12.
- Akubor PI, Badifu GIO. 2004. Chemical composition, functional properties and baking potential of African breadfruit kernel and wheat flour blends. *Int. J. Food Sci. Technol.* 39:223-9.
- Akusu MO, Wordu GO. 2013. Nutritional and sensory evaluation of EPITI (IKWOKE) prepared from maize/soybean flour blends. *Int. J. Sci. Emerg. Technol.* 6(3):230-5.
- AOAC. 1990. Official methods of analysis, 15th Eds, Washington, 774 p.
- Aremu MO, Olonisakin A, Atolaye BO, Ogbu CF. 2007. Some nutritional composition and functional properties of *Prosopis Africana*. *Bangladesh J. Sci. Ind. Res.* 42 (3):269-80.
- Atwater WO, Benedict FG. 1902. Experiments on the metabolism of matter and energy in the human body. Bulletin N° 109, Washington DC.
- Bekoye BM, Akanvou R. 2005. Bien cultiver le mil en Côte d'Ivoire. Centre National de Recherche Agronomique (CNRA); 4 p.
- Demaison L, Moreau D. 2002. Dietary n-3 polyunsaturated fatty acids and coronary heart disease-related mortality: a possible mechanism of action. *Cell. Molecul. Life Sci.* 5:463-77.
- Duc G, Mignolet C, Carrouée B, Huyghe C. 2010. Importance économique passée et présente des légumineuses : rôle historique dans les assolements et facteurs d'évolution. *Innov. Agronom.* 11:1-24.
- El-Adawy TA. 2000. Functional properties and nutritional quality of acetylated and succinylated mung bean protein isolate. *Food Chem.* 70 (1):83-91.
- FAO. 1995. Le sorgho et les mils dans l'alimentation humaine. <http://www.fao.org/docrep/t0818f/t0818f01.htm> (15/01/2016)
- FAO. 2009. La situation mondiale de l'alimentation et l'agriculture. <http://www.fao.org/datalog/Inter.htm> (15/01/2016).
- Folch J, Lees M, Stanley GHS. 1957. A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* 172:497-509.
- Kalimbira AA, Mtimuni BM, Mtimuni JP. 2004. Effect of incorporating legumes on nutritive value of cassava-based complementary foods. *J. Agric. Environ. Sci. Technol.* 2:13-21.
- Lin MJY, Humbert ES, Sosulski FW. 1974. Certain functional properties of sunflower meal products. *J. Food Sci.* 39:368-71.
- Mouquet C, Bruyeron O, Trèche S. 1998. Les farines infantiles : Caractéristiques d'une bonne farine infantile. *Bull. Res. TPA* 15:11.
- Moure A, Sineiro J, Dominiguez H, Parajo J.C. 2006. Functionality of oil seed protein product: a review. *Food Res. Int.* 38:945-63.
- Narayana K, Narasimma RNMS. 1982. Functional properties of raw and heat processed winged bean flour. *J. Food Sci.* 47:1534-38.
- Njintang NY, Mbofung CMF, Waldron KW. 2001. *In vitro* protein digestibility and physicochemical properties of dry red bean (*Phaseolus vulgaris*) flour: effect of processing and incorporation of soybean and cowpea flour. *J. Agri. Food Chem.* 49:2465-71.
- Phillips RD, Chinnan MS, Branch AL, Miller J, Mcwatters KH. 1988. Effects of pre-treatment on functional and nutritional properties of cowpea meal. *J. Food Sci.* 3:805-9.
- Sakai WS. 1979. Aroid root crop In: *Tropical Foods*, Chem. Nutr., New York 1, p 268.
- Sanogo M. 1994. La production artisanale de farines infantiles. Editions du GRET, Paris, 79 p.
- Shad MA, Nawaz H, Noor M, Ahmad HB, Hussain M, Choudhry MA. 2013. Functional properties of maize flour and its blends with wheat flour: optimization of preparation conditions by response surface methodology. *Pakistan J. Botany* 45(6):2027-35.
- Siddiq M, Rav R, Harte JB, Dolan KD. 2010. Physical and functional characteristics of selected dry bean (*Phaseolus vulgaris* (L.) flours. *J. Food Sci. Technol.* 43:232-7.
- Soro S, Konan G, Elleingand E, N'guessan D, Koffi E. 2013. Formulation d'aliments infantiles à base de farines d'igname enrichies au soja. *Afr. J. Food Agric. Nutr. Dev.* 13(5):8313-39.
- Sosulski FW. 1962. The centrifuge method for determining flour absorption in hard red spring wheat. *Cereal Chem.* 39:344-50.
- Wolf WJ. 1970. Soybean proteins: their functional, chemical and physical properties. *J. Agric. Food Chem.* 18:965-9.
- Yasumatsu K, Sawada K, Moritaka S, Misaki M, Tuda J, Wada T, Ishii K. 1972. Whipping and emulsifying properties of soybean products. *Agri. Biol. Chem.* 36:716-27.
- Zannou-Tchoko JV, Ahui-Bitty BL, Kouame K, Bouaffou MGK, Dally T. 2011. Utilisation de la farine de maïs germe source d'alpha amylases pour augmenter la densité énergétique de bouillies de sevrage à base de manioc et son dérivé *attiéké*. *J. Appl. Biosci.* 37:2477-84.