



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

ASIAN JOURNAL OF
SCIENCE AND TECHNOLOGY

Asian Journal of Science and Technology
Vol. 5, Issue 3, pp. 198-203, March, 2014

RESEARCH ARTICLE

THEORETICAL STUDY OF THE EFFECTS OF SOLVENTS ON GLOBAL PROPERTIES OF DDQ

¹*Srivastava, K. K., ²Shubha Srivastava, ³Md. Tanweer Alam and ⁴Rituraj

^{1,3,4}Department of Chemistry, Vinoba Bhave University, Hazaribagh, India

²Department of Chemistry, KBW College, Hazaribagh, India

ARTICLE INFO

Article History:

Received 07th December, 2013

Received in revised form

11th January, 2014

Accepted 28th February, 2014

Published online 25th March, 2014

ABSTRACT

DDQ is widely used as a π - acceptor for the preparation of organic charge transfer. In this paper the effect of ten solvents on the ground state of DDQ has been reported. DFT calculations have been done on the Schrodinger software and the effect of solvents have been theoretically calculated with the help of Poisson-Boltzmann solver. The solvation energy, chemical potential, hardness, electrophilicity, HOMO-LUMO gap and the picture of the HOMO and LUMO of DDQ in the ground state in the solvents have been reported.

Key words:

DFT, Ground state,

DDQ, Energy components,

HOMO, LUMO,

Solvation energy,

Dielectric constants.

Copyright © 2014 Srivastava et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Physical and chemical property of a molecule depends on the structure and the various kinds of energies of the molecule. Chemical reaction of a molecule in solution is affected by the nature of the solvent; solvent affects not only the energies of HOMO and LUMO of the molecule, but also other kinds of energies. Energy of a molecule may be considered to have various energy components such as reaction field energy, total zero-electron terms, Nuclear-nuclear, Nuclear-solvent, total one-electron terms, Electron-nuclear, Electron-solvent, Kinetic, total two-electron terms, Electronic energy, total quantum mech. energy, Gas phase energy, Solution phase energy, total solute energy, total solvent energy, Solute cavity energy, Reorganization energy, Solvation energy total internal energy, total enthalpy, total Gibbs free energy, and zero point energy. DDQ is a Bright yellow solid, It is used in preparation of Charge transfer complexes (Foster Roy, 1969; Gutman and Lyons, 1969; Kandana and Pohl, 1968; Slifkin, 1971; LamisShahada et al., 2009; Vinod Kumar et al., 2000; Becker, 1965; Subhan et al., 2000; Refat Moamen and El-Metwally Nashwa, 2011; Liang-YeuChen et al., 2010) of various utilities. DDQ is also one of the most versatile reagent because of its high oxidant ability and relative stability compared to others. Keeping in view the utility of DDQ various kinds of energies of DDQ in the ground state in gaseous phase and in different kinds of solvents have been theoretically calculated in this paper.

*Corresponding author: Srivastava, K. K., Department of Chemistry, Vinoba Bhave University, Hazaribagh, India

Computational methods

The initial structure of DDQ was built with Chem-Draw ultra8.0 and the structure was optimized on Chem3D ultra 8.0. The structure was exported to Maestro 9.3 of Schrodinger 2012 version. The optimization of the structure was done on the Jaguar panel of the Maestro 9. The DFT-B3LYP method of theory was chosen. 6-31g^{##} basis set was selected and 255 basis functions were created for calculation. The molecule was assigned net zero charge and singlet multiplicity. In the solvent menu of the jaguar panel PBF solver was used for optimization of the structure in both the gaseous and solution phase. The optimization the gaseous state and in the different solutions were done in ground state of the molecule.

Geometry optimization

for perform a geometry optimization one needs to guess at the geometry and the direction in which to search, a set of co-ordinates to optimize, and some criteria for when to optimization is complete. The search direction is obtained from the gradient of the energy and the initial Hessian. An initial Hessian (second derivative matrix or force constant matrix) and the gradient are used to define search direction that should result in lowering of energy. The choice if co-ordinate systems have a substantial impact on the convergence of the optimization. The ideal set of Co-ordinate is one in which the energy change along each co-ordinate is maximized, and the coupling between co-ordinates is minimized. Jaguar chooses the coordinate system by default. It has two options

Cartesian and z-matrix that produces an efficient optimization requires an understanding of the coupling between simple internal co-ordinates. For optimization to minimum energy structures, the convergence criterion for SCF calculation is chosen to assure accurate analyses gradients. For these jobs, a wave function is considered converged when the root mean square (RMS) change in density matrix element is less than the RMS density matrix element change criterion, whose default value is 5.0×10^{-6} . The geometry is considered to have converged when the energy of successive geometries and the elements of analyze gradients of the energy and the displacement has met convergence criteria. For optimization in solution, the default criteria are multiplied by a factor of three, and a higher priority is given to the energy convergence criterion. Thus if the energy change criterion is met before the gradient and displacement criteria have been met, the geometry is considered converged. The optimized geometry may not have a local minimization energy i.e it may have reside on a saddle. To know whether it is global minimization we look for the value of vibrational frequencies. If all the vibrational frequencies are real (i.e+ve) then it represents global minimum, but if any of the vibrational frequencies is negative (i.e imaginary) then it is local minimum.

Performing a solvation calculation

It involves several iterations in which the wave functions for the molecule in the gas phase are calculated. The program *ch* performs electrostatic potential fitting, which represents the wave function as a set of point charges on the atomic centers. The interactions between the molecule and the solvent are evaluated by Jaguar's Poisson-Boltzmann solver (Tannor et al., 1994; Marten et al., 1996), which fits the field produced by the solvent dielectric continuum to another set of point charges. These charges are passed back to *scf*, which performs a new calculation of the wave function for the molecule in the field produced by the solvent point charges. Electrostatic potential fitting is performed on the new wave function, the solvent-molecule interactions are reevaluated by the Poisson-Boltzmann solver, and so on, until the solvation free energy for the molecule converges.

For solvation calculations on neutral systems in water the program *pre* evaluates the Lewis dot structure for the molecule or system and assigns atomic van der Waals radii accordingly. These van der Waals radii are used to form the boundary between the solvent dielectric continuum and the solute molecule. The Lewis dot structure and van der Waals radii information both appear in the output from the program *pre*. The radii are listed under the heading "vdw2" in the table of atomic information below the listing of non-default options. After the *pre* output, the usual output appears for the first, gas-phase calculation, except that the energy breakdown for the *scf* output also describes the electron-nuclear and kinetic contributions to the total one-electron terms in the energy, as well as the virial ratio $-V/T$, where V is the potential energy and T is the kinetic energy. This ratio should be -2 if the calculation satisfies the virial theorem. After the first *scf* output, the output from the first run of the program *ch* appears. Since performing a solvation calculation enables electrostatic potential fitting to atomic centers, the usual output for that option is included every time output from the program *ch* appears in the output file. The post program writes out the necessary input files for the Poisson-Boltzmann solver; this

step is noted in the output file. The next output section comes from the Poisson-Boltzmann solver. The output includes information on the area (in \AA^2) of the molecular surface formed from the intersection of spheres with the van der Waals radii centered on the various atoms; the reaction field energy in kT (where $T = 298$ K), which is the energy of the interaction of the atom-centered charges with the solvent; the solvent-accessible surface area (in \AA^2), which reflects the surface formed from the points whose closest distance from the molecular surface is equal to the probe radius of the solvent; and the cavity energy in kT , which is computed to be the solvation energy of a nonpolar solute whose size and shape are the same as those of the actual solute molecule. The output from the program *solv* follows the Poisson-Boltzmann solver results, giving the number of point charges provided by the solver to model the solvent, the sum of the surface charges, the nuclear repulsion energy already calculated by Jaguar, the nuclear-point charge energy representing the energy of interaction between the molecule's nuclei and the solvent point charges, and the point-charge repulsion energy, which is calculated but not used because it is irrelevant to the desired solvation results. After this output, the output for the second solvation iteration begins.

The output from *scf* comes first, giving the results for the molecule-and-solvent-point-charges system. Total quantum mech. energy corresponds to the final energy from the *scf* energy table for that iteration, and includes the entire energies for the molecule-solvent interactions. The output next includes the gas phase and the solution phase energies for the molecule, since these terms are, of course, necessary for solvation energy calculations. The first solution phase energy component is the total solute energy, which includes the nuclear-nuclear, electron nuclear, kinetic, and two-electron terms, but no terms involving the solvent directly. The second component of the solution phase energy is the total solvent energy, which is computed as half of the total of the nuclear-solvent and electron-solvent terms, since some of its effect has already changed the solute energy. Third, a solute cavity term, which computes the solvation energy of a nonpolar solute of identical size and shape to the actual solute molecule, as described in reference (Tannor et al., 1994), is included. This is only done for water as solvent. The last solution phase energy component (shown only if it is nonzero) is term (T), the first shell correction factor, which depends on the functional groups in the molecule, with atoms near the surface contributing most heavily. Finally, the list ends with the reorganization energy and the solvation energy. The reorganization energy is the difference between the total solute energy and the gas phase energy, and does not explicitly contain solvent terms. The final solvation energy is calculated as the solution phase energy described above minus the gas phase energy. The solvation energy is listed in Hartrees and in kcal/mol,

Chemical potential (μ) (Parr et al., 1983)

HOMO as ionization energy (IE) and LUMO as electron affinity (EA) have been used for calculating the electronic chemical potential (μ) which is half of the energy of HOMO and LUMO $\mu = (E_{\text{HOMO}} + E_{\text{LUMO}})/2$

Hardness (η) (Pearson, 1991)

The hardness (h) as half of the gap energy of HOMO and LUMO has been calculated using the following equation $\text{Gap} = E_{\text{HOMO}} - E_{\text{LUMO}}$ $\eta = \text{Gap}/2$

Electrophilicity (ω) (Parr and Yang, 1989)

The electrophilicity (ω) has been calculated using equation $\omega = \mu^2/2\eta$

Reaction field energy (in KT)

This gives us the energy of the interactions of atom centred charges with the solvent; Solvent accessible surface area (SASA in Å^2) reflects the surface formed from the points whose closest distance from the molecular surface is equal to the probe radius of the solvent.

Cavity energy (in KT)

This is solvation energy of a non-polar solute whose size and shape are the same as those of actual solute molecule.

Quantum mechanical energy

This term corresponds to the entire energies for the molecule solvent interaction and is equal to the sum of total zero electron terms and electronic energy.

Reorganisation energy

This is the difference between the total solute energy and the gas phase energy, and does not explicitly contain solvent terms.

RESULTS AND DISCUSSION**Solvent parameters**

Table-1 summarizes the solvent parameters such as dielectric constants, molecular weight, density and polarity of the solvents used for the present theoretical study by Poisson-Boltzmann solver. In table-3 the energy components calculated by DFT method on Jaguar panel of the Maestro 9.3 with 6-31g^{##} basis set utilizing 255 basis functions for 1,3,5-trinitrobenzene in the ground state have been incorporated. The pictures of HOMO and LUMO of DDQ in gaseous state and in various solvents have been shown in fig.1

Table 1. Physical parameters of various solvents

Solvents	M.W	Density	Dielectric constant	Probe radius
	g/mol	g/ml		Å ⁰
1.Acetonitrile	37.5	0.777	37.5	2.19
2.Benzene	78.12	0.87865	2.284	2.6
3.Carbondetrachloride	153.82	1.594	2.238	2.67
4.Chloroform	119.38	1.4832	4.806	2.52
5.Cyclohexane	84.16	0.77855	2.023	2.78
6.Dichloromethane	84.93	1.3266	8.93	2.33
7.DMF	73.09	0.944	36.7	2.49
8.methanol	32.04	0.7914	33.62	2
9.THF	72.11	0.8892	7.6	2.52
10.Water	18.02	0.99823	80.37	1.4

Table 2. Values of HOMO-LUMO energy, μ , η , ω of DDQ calculated by DFT -B3LYP/6-31G- level

Solvents	HOMO	LUMO	Gap	$\mu = E_{\text{HOMO}} + E_{\text{LUMO}}/2$	$\eta = (E_{\text{LUMO}} - E_{\text{HOMO}})/2$	Electrophilicity $\omega = \mu^2/2\eta$
HOMO, LUMO energy in eV						
acetonitrile	-8.018	-4.817	-3.201	-6.417	1.601	0.045
benzene	-8.233	-4.984	-3.249	-6.608	1.624	0.048
carbontetrachloride	-8.237	-4.987	-3.250	-6.612	1.625	0.048
chloroform	-8.128	-4.908	-3.221	-6.518	1.610	0.046
cyclohexane	-8.255	-5.000	-3.255	-6.628	1.627	0.048
dichloromethane	-8.073	-4.863	-3.211	-6.468	1.605	0.045
dmf	-8.019	-4.820	-3.198	-6.419	1.599	0.044
methanol	-8.020	-4.816	-3.204	-6.418	1.602	0.045
THF	-8.085	-4.874	-3.211	-6.480	1.606	0.046
Water	-8.035	-4.856	-3.179	-6.446	1.590	0.045

An electron acceptor represents the ability to obtain an electron in the LUMO and HOMO represents the ability to donate electron. The ($E_{\text{HOMO}} - E_{\text{LUMO}}$) gap is an important scale of stability (Johnson et al., 2010) and compounds with large ($E_{\text{HOMO}} - E_{\text{LUMO}}$) gap value tend to have higher stability. The perusal of the table-2 indicates the stability of DDQ increases in the solvents in the ground state in the order;

water > dmf > acetonitrile > methanol > THF = dichloromethane > chloroform > benzene > carbontetrachloride > cyclohexane

The plot of the energy gap between HOMO and LUMO versus dielectric constant of solvents in ground state have been shown in the fig 2. The dependence of the energy gap (y) on dielectric constant (x) in ground state follows the equation $y = 1E-08x^5 - 2E-06x^4 + 0.0001x^3 - 0.0031x^2 + 0.0288x - 3.3007$ ($R^2 = 0.999$).

The DDQ molecule has been found to be stabilized in the ground state in *water > dmf > acetonitrile > methanol > THF = dichloromethane > chloroform > benzene > carbontetrachloride > cyclohexane* r. Therefore, if it is desired to stabilize DDQ in the ground state then out of ten solvents studied water is the best.

The 3D plots of HOMO-LUMO and dielectric constant shown in figure 7. reveals that LUMO of DDQ is more effected than HOMO in both the ground state by change in the dielectric constant of the solvent.

The chemical potentials (μ) of DDQ in the ground state increases in the order;

acetonitrile > methanol > dmf > water > dichloromethane > THF > chloroform > benzene > carbontetrachloride > cyclohexane.

The plot of the chemical potential versus dielectric constant of solvents in ground state have been shown in the fig3. The dependence of the chemical potential (y) on dielectric constant(x) follows the equation $y = 3E-08x^5 - 6E-06x^4 + 0.0003x^3 - 0.0079x^2 + 0.0804x - 6.757$ ($R^2 = 0.9992$).

The DDQ molecule has been found to possess higher chemical potential in the ground state in benzene, chloroform, gas, methanol, water in the order;

acetonitrile > methanol > dmf > water > dichloromethane > THF > chloroform > benzene > carbon-tetrachloride > cyclohexane.

Therefore, if it is desired to have highest chemical potential, DDQ in the ground state, then out of ten solvents studied acetonitrile is the best.

Table 3. Values of energy components of DDQ in gaseous state and various solvents in ground state calculated by DFT -B3LYP/6-31G- level

Ground state											
Energy components, in eV	Gas-phase	Acetonitrile	Benzene	Carbon tetrachloride	Chloroform	Cyclohexane	Dichloro methane	dmf	methanol	THF	Water
(A)Total zero electron terms		26343.39	26373.61	26363.60	26354.28	26365.43	26349.23	26343.67	26343.46	26350.48	26350.86
(B)Nuclear-nuclear	26373.15	26396.08	26395.37	26384.88	26390.57	26384.20	26393.57	26396.00	26396.28	26392.88	26395.79
(C)Nuclear-solvent		-52.69	-21.75	-21.28	-36.29	-18.77	-44.34	-52.33	-52.82	-42.40	-44.92
(E)Total one electron terms	-107805.96	-107772.93	-107838.06	-107797.58	-107786.50	-107799.94	-107780.46	-107773.27	-107773.12	-107781.98	-107786.0
(F)Electron-nuclear		-148017.19	-148067.61	-148011.86	-148015.11	-148011.85	-148016.75	-148017.13	-148017.48	-148016.43	-148022.9
(G)Electron-solvent		51.36	21.26	20.80	35.43	18.36	43.26	51.01	51.48	41.37	43.87
(H)Kinetic		40192.91	40193.51	40193.48	40193.19	40193.55	40193.04	40192.86	40192.88	40193.08	40193.01
(I)Total two electron terms	41021.92	41017.47	41023.17	41022.64	41020.54	41023.23	41019.36	41017.53	41017.59	41019.68	41023.30
(L)Electronic energy (E+I)	-66784.04	-66755.46	-66775.27	66774.94	-66765.96	-66776.71	-66761.10	-66755.73	-66755.53	44574.81	44578.44
(N)Total quantum mechanical energy (A+L)	-40410.89	-40412.07	-40411.35	-40411.34	-40411.68	-40411.28	-40411.87	-40412.06	-40412.08	-40411.82	-40411.84
(O)Gas phase energy		-40410.89	-40410.89	-40410.89	-40410.89	-40410.89	-40410.89	-40410.89	-40410.89	-40410.89	-40410.89
(P)Solution phase energy (Q+R+S)		-40411.40	-40411.11	-40411.10	-40411.25	-40411.08	-40411.32	-40411.40	-40411.41	-40411.31	-40411.19
(Q)Total solute energy (N-C-G)		-40410.74	-40410.86	-40410.86	-40410.82	-40410.87	-40410.78	-40410.74	-40410.74	-40410.79	-40410.79
(R)Total solvent energy C/2+G/2)		-0.67	-0.24	-0.24	-0.43	-0.21	-0.54	-0.66	-0.67	-0.52	-0.53
(S)Solute cavity energy	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
(U)Reorganization energy (Q-O)		0.15	0.03	0.03	0.07	0.02	0.11	0.15	0.15	0.10	0.10
(V)solvation energy (P-O) (kJ/mol)		-11.86	-4.97	-4.86	-8.24	-4.30	-10.00	-11.78	-11.89	-9.60	-6.81

The hardness(η) of DDQ increases in the ground state in the following order;

Cyclohexane>carbontetrachloride> benzene>chloroform>THF>dichloromethane> methanol>acetonitrile>dmf>water.

The plot of hardness versus dielectric constant of solvents in the ground state have been shown in the fig4. The dependence of hardness (y) on dielectric constant(x) follows $y = -7E-09x^5 + 1E-06x^4 - 7E-05x^3 + 0.0016x^2 - 0.0144x + 1.6504$ ($R^2 = 0.999$)

The DDQ molecule has been found to be hardest in cyclohexane in the ground state. Therefore, if it is desired to increased hardness of DDQ to largest extent in the ground state then out of ten solvents studied cyclohexane is the best.

The electrophilicity (ω) of DDQ increases in ground state in the following order

Benzene=carbontetrachloride=cyclohexane>chloroform=THF>acetonitrile=dichloromethane=methanol=water>dmf

The plot of electrophilicity (y) versus dielectric constant(x) of solvents in ground state have been shown in the fig5. The dependence of the electrophilicity on dielectric constant follows the $y = 2E-11x^6 - 5E-09x^5 + 3E-07x^4 - 1E-05x^3 + 0.0002x^2 - 0.0018x + 0.0511$ ($R^2 = 1.0002$) in the ground.

The DDQ molecule has been found to possess high electrophilicity in the ground in benzene, carbontetrachloride and cyclohexane. Therefore, if it is desired to increase electrophilicity of DDQ to larger extent in the ground state, then out of ten solvents studied benzene, carbontetrachloride and cyclohexane are the best.

The Solvation energy of DDQ in the ground state are in the following order; methanol> acetonitrile>dmf> dichloromethane> THF> chloroform> water> benzene>carbontetrachloride> cyclohexane.

The plot of the solvation energy versus dielectric constant of solvents in ground state have been shown in the fig.6. The dependence of the solvation energy(y) on dielectric constant(x) follows

$$y = -1E-06x^5 + 0.0002x^4 - 0.0119x^3 + 0.2871x^2 - 2.9087x + 0.3833$$
 ($R^2 = 0.9992$)

Thus, it was found that DDQ is most highly solvated in the ground state in methanol than other studied solvents.

State	HOMO	LUMO
Gaseous state		
cyclohexane		
carbontetrchloride		
benzene		
chloroform		
THF		
dichloromethane		
methanol		
DMF		
acetonitrile		
water		

Figure 1. Picture of HOMO-LUMO of DDQ in the ground state in gaseous and ten various solvents




Figure 2. Effect of dielectric contant on the HOMO-LUMO gap of DDQ in the GS




Figure 3. Effect of dielectric contant on the chemical potential of DDQ in the GS




Figure 4. Effect of dielectric contant on the hardness DDQ in the GS




Figure 5. Effect of dielectric contant on the electrophilicity of DDQ in the GS




Figure 6. Effect of dielectric constant on the solvation of DDQ in the GS




Figure 7. Figure 3D plot of HOMO-LUMO-dielectric constant to study the effect on HOMO and LUMO of DDQ with the change in the dielectric constant of solvents in ground state

Conclusion

The present study on solvent effect on the energy components of DDQ in ground state by ten different solvents has led us to conclude it is highly solvated in methanol while lowest in cyclohexane. It has been found that DDQ is most hard, electrophilic and chemical potential in cyclohexane, benzene (also in cyclohexane, and carbon-tetrachloride, and acetonitrile respectively while least in water, dmf and cyclohexane respectively.

Acknowledgment

We are thankful to Will Richard, Raghu Rangaswamy, Vinod Dewarjee for providing the Shrodinger2012 software.

REFERENCES

- Foster Roy, Organic charge transfer complexes, Academic Press, New York (1969).
 Gutman F and L E Lyons, Organic semiconductors, Lohm Willey and Sons, Inc. New York (1969).
 Kandana S and H A Pohl, in organic semiconducting polymers Ed. J E Katon, Marcel Dekker Inc., New York (1968) p. 86.

Slifkin, M A Charge transfer interactions of biomolecules, Academic Press, New York, (1971).

Lamis Shahada a, Adel Mostafa b, El-Metwally Nour a, Hassan S. Bazzi Journal of Molecular Structure Synthesis, spectroscopic, thermal and structural investigations of charge-transfer complexes of 4,40-trimethylenedipiperidine with chloranil, TBCHD, DDQ, TCNQ and iodine Journal of Molecular Structure 933 (2009) 1–7

VINOD KUMAR, T, T VEERAAIAH and G VENKATESHWARLU, Molecular complexes of phenols with DDQ Proc. Indian Acad. Sci. (Chem. Sci.), Vol. 112, No. 2, April 2000, pp. 119–125 *Indian Academy of Sciences*

Becker H D 1965 *J. Org. Chem.* 30 982

Hewgill F R and Howie G B 1978 *Aust. J. Chem.* 31 907

SUBHAN I, M. S., N. K. BHATT I, M. MOHAMMAD, A. Y. KHAN Turk J Chem 24 (2000), 223 {230.cTUBITAK

REFAT Moamen & El-METWALLY Nashwa M July 2011 Vol.56 No.19: 1993–2000doi: 10.1007/s11434-011-4525-9Refat M S, et al. Chinese Sci Bull July (2011) Vol.56 No.19 Investigation of charge transfer complexes formed between 3,3'-dimethylbenzidine (o-toluidine) donor and DDQ, p-chloranil and TCNQ as π -acceptors

Liang-YeuChen, a, bSie-RongLi, a Po-Yuan Chen, a Ho-Chiang Chang, a Tzu-Pin Wang, alan-LihTsai, b and Eng-Chi Wang,* 2,3-Dichloro-5,6-dicyanobenzoquinone (DDQ) mediated oxidation dehydrogenation of 2-aryl-3,4-dihydro-2H-benzopyrans: synthesis of 2-arylbenzopyran-4-ones ARKIVOC 2010 (xi) 64-76

Tannor, D. J.; Marten, B.; Murphy, R.; Friesner, R. A.; Sitkoff, D.; Nicholls, A.; Ringnalda, M.; Goddard, W. A., III; Honig, B. Accurate First Principles Calculation of Molecular Charge Distributions and Solvation Energies from Ab Initio Quantum Mechanics and Continuum Dielectric Theory. *J. Am. Chem. Soc.* 1994, 116, 11875

Marten, B.; Kim, K.; Cortis, C.; Friesner, R. A.; Murphy, R. B.; Ringnalda, M. N.; Sitkoff, D.; Honig, B. New Model for Calculation of Solvation Free Energies: Correction of Self-Consistent Reaction Field Continuum Dielectric Theory for Short-Range Hydrogen-Bonding Effects. *J. Phys. Chem.* 1996, 100, 11775.

Parr, R. G.; Pearson, R. G. *J. Am. Chem. Soc.*, 1983, 105, 7512.

Pearson, RG 1991, Density functional theory-electro negativity and hardness chemtracts- *Inorg.chem.* 3:317-333
 Parr, R.G.; Yang, W. Chemical potential derivatives. In *Density-Functional Theory of Atoms and Molecules*, 1st ed.; Oxford University Press: New York, NY, USA, 1989; pp. 87–95.

Johnson, E. R.; Keinan, S.; Mori-Sanchez, P.; Contreras-Garcia, J.; Cohen, A. J.; Wang, W. *J. Am. Chem. Soc.* 2010, 132, 6498.
